

# From Sparse to Dense Functional Data in High Dimensions: Revisiting Phase Transitions from a Non-Asymptotic Perspective

**Shaojun Guo**

*Institute of Statistics and Big Data  
Renmin University of China  
Beijing 100872, China*

SJGUO@RUC.EDU.CN

**Dong Li**

*Department of Statistics and Data Science  
Tsinghua University  
Beijing 100084, China*

MALIDONG@TSINGHUA.EDU.CN

**Xinghao Qiao\***

*Faculty of Business and Economics,  
The University of Hong Kong  
Hong Kong, China*

XINGHAOQ@HKU.HK

**Yizhu Wang**

*Department of Mathematical Sciences  
Tsinghua University  
Beijing 100084, China*

WANGYIZH20@MAILS.TSINGHUA.EDU.CN

**Editor:** Genevera Allen

## Abstract

Nonparametric estimation of the mean and covariance functions is ubiquitous in functional data analysis and local linear smoothing techniques are most frequently used. Zhang and Wang (2016) explored different types of asymptotic properties of the estimation, which reveal interesting phase transition phenomena based on the relative order of the average sampling frequency per subject  $T$  to the number of subjects  $n$ , partitioning the data into three categories: “sparse”, “semi-dense”, and “ultra-dense”. In an increasingly available high-dimensional scenario, where the number of functional variables  $p$  is large in relation to  $n$ , we revisit this open problem from a non-asymptotic perspective by deriving comprehensive concentration inequalities for the local linear smoothers. Besides being of interest by themselves, our non-asymptotic results lead to elementwise maximum rates of  $L_2$  convergence and uniform convergence serving as a fundamentally important tool for further convergence analysis when  $p$  grows exponentially with  $n$  and possibly  $T$ . With the presence of extra  $\log p$  terms to account for the high-dimensional effect, we then investigate the scaled phase transitions and the corresponding elementwise maximum rates from sparse to semi-dense to ultra-dense functional data in high dimensions. We also discuss a couple of applications of our theoretical results. Finally, numerical studies are carried out to confirm the established theoretical properties.

---

\*. The authors' names are sorted alphabetically, and the corresponding author is Xinghao Qiao.

**Keywords:** concentration inequalities, high-dimensional partially observed functional data, elementwise maximum rates, local linear smoother, mean and covariance functions

## 1. Introduction

A fundamental issue in functional data analysis is the nonparametric estimation of the mean and covariance functions based on discretely sampled and noisy curves. Despite being of interest by themselves, the estimated quantities serve as building blocks for dimension reduction and subsequent modeling of functional data, such as functional principal component analysis (FPCA) (Yao et al., 2005a; Li and Hsing, 2010) and functional linear regression (Yao et al., 2005b; Chen et al., 2022). Among candidate nonparametric smoothers, we focus on the most commonly-adopted local linear smoothing method due to its simplicity and attractive local and boundary correction properties.

In a typical functional data setting, we have  $n$  random curves, representing  $n$  subjects, observed with errors, at  $T_i$  randomly sampled time points for the  $i$ th subject. The sampling frequency  $T_i$  plays a pivotal role in the estimation, as it may affect the choice of the estimation procedure. The literature can be loosely divided into two categories. The first category corresponds to dense functional data, where  $T_i$ 's are larger than some order of  $n$ . A conventional approach to handle such data implements nonparametric smoothing to the observations from each subject to eliminate the noise, thus reconstructing each individual curve before subsequent analysis (Zhang and Chen, 2007). The second category referred to as sparse functional data, accords with bounded  $T_i$ 's. Under such a scenario, the pre-smoothing step is no longer applicable, an alternative pooling strategy considers pooling the data from all subjects to build strength across all observations (Yao et al., 2005a; Li and Hsing, 2010). More recently, Zhang and Wang (2016) provided a comprehensive analysis of phase transitions and the associated rates of convergence for three types of asymptotic properties: local asymptotic normality,  $L_2$  convergence, and uniform convergence. They proposed to further partition dense functional data into new categories: “semi-dense” and “ultra-dense”, depending on whether the root- $n$  rate is achieved with negligible asymptotic bias or not. However, these aforementioned asymptotic results are only suitable for handling univariate or low-dimensional multivariate functional data.

With recent advances in data collection technology, high-dimensional functional data sets become increasingly available. Examples include time-course gene expression data, and electroencephalography and functional magnetic resonance imaging data, where signals are measured over time at a large number of regions of interest (Zhu et al., 2016; Li and Solea, 2018; Zapata et al., 2022; Fang et al., 2024). Those data can be represented as a  $p$ -vector of random functions  $\mathbf{X}_i(\cdot) = \{X_{i1}(\cdot), \dots, X_{ip}(\cdot)\}^T$  for  $i = 1, \dots, n$  defined on a compact set  $\mathcal{U}$ , with the  $p$ -vector of mean functions  $\boldsymbol{\mu}(\cdot) = \{\mu_1(\cdot), \dots, \mu_p(\cdot)\}^T = \mathbb{E}\{\mathbf{X}(\cdot)\}$  and the  $(p \times p)$ -matrix of marginal- and cross-covariance functions

$$\boldsymbol{\Sigma}(u, v) = \{\Sigma_{jk}(u, v)\}_{p \times p}, \quad \Sigma_{jk}(u, v) = \text{cov}\{X_{ij}(u), X_{ik}(v)\}. \quad (1)$$

In a high-dimensional regime, the dimension  $p$  can be diverging with, or even larger than, the number of subjects  $n$ . In practice, each  $X_{ij}(\cdot)$  is observed subject to error contamination at  $T_{ij}$  random time points. See (4) below.

The estimation of mean and covariance functions for high-dimensional functional data is not only interesting at its own right but also plays a foundational role as a critical starting point for subsequent analysis. Due to the infinite-dimensionality of functional data, it is common practice to truncate the basis expansion of each function  $X_{ij}(\cdot)$  at a finite level, using either data-driven basis expansion via FPCA or pre-fixed basis expansion. This allows for the subsequent development of regularized methods for the sparse estimation to address the high-dimensionality based on estimated FPC scores or estimated basis coefficients. Importantly, the estimation of mean and covariance functions are implicitly involved in this procedure. Applications of this type of approach to estimate sparse high-dimensional functional models include functional graphical models (Qiao et al., 2019; Zhao et al., 2022; Solea and Li, 2022; Zapata et al., 2022; Lee et al., 2023), functional additive regressions (Fan et al., 2014, 2015; Kong et al., 2016; Luo and Qi, 2017; Xue and Yao, 2021), sparse FPCA (Hu and Yao, 2022) and functional linear discriminant analysis (Xue et al., 2024). Another line of applications considers the sparsity-induced estimation of the covariance matrix function without any basis expansion, such as Fang et al. (2024); Li et al. (2023) and Leng et al. (2024). Both types of applications call for the investigation of non-asymptotic properties of the componentwise mean and covariance estimators, which serves as the motivation of our paper when dealing with the practical scenario of partially observed curves in high dimensions. See Section 4 for details of some applications.

Within the high-dimensional statistical learning framework, it is essential to conduct non-asymptotic analysis of the estimators by developing concentration inequalities under a given performance metric, which can lead to probabilistic error bounds in the elementwise maximum norm as a function of  $n$ ,  $p$ , and possibly  $T_{ij}$ 's (depending whether they are diverging or bounded) under our setup. Existing literature has mainly focused on fully observed functional data, based on which concentration inequalities for the estimated covariance functions were established in Qiao et al. (2019) and Zapata et al. (2022). In practical scenarios where curves are partially observed with errors, addressing dense functional data is achievable by applying the pre-smoothing technique to observations from each  $i, j$  (Kong et al., 2016). Alternatively, a unified pooling-type local linear smoothing approach can be employed for estimating the mean functions  $\mu_j(\cdot)$ 's and marginal- and (or) cross-covariance functions  $\Sigma_{jk}(\cdot, \cdot)$ 's across  $j, k$  to handle both sparsely and densely observed functional data (Qiao et al., 2020; Lee et al., 2023; Fang et al., 2024). Although such nonparametric smoothing approach suffers from high computational cost when  $p$  is large, it can be substantially accelerated in a common practical scenario where each  $X_{ij}(\cdot)$  is observed at the same set of points across  $j \in [p]$  especially with the aid of linear binning (Fan and Marron, 1994), resulting in an efficient estimation procedure. Moreover, those commonly-adopted FPCA-based methods only necessitate the estimation of marginal- instead of cross-covariance functions across  $j \in [p]$  (Qiao et al., 2019; Solea and Li, 2022), and can be easily paralleled for fast computation. See Remark 1.

On the theory side, this approach entails dealing with the second-order  $U$ -statistics with complex dependence structures, posing a technically challenging task. Qiao et al. (2020) made the first attempt to derive some sub-optimal concentration inequalities for local linear smoothers of marginal-covariance functions  $\hat{\Sigma}_{jj}(\cdot, \cdot)$ 's, albeit under a restrictive finite-dimensional setting. Lee et al. (2023) established the convergence of their proposed estimation of conditional functional graphical models under the assumption of elementwise

maximum rate for the covariance smoothers:

$$\max_{1 \leq j, k \leq p} \|\widehat{\Sigma}_{jk} - \Sigma_{jk}\|_{\mathcal{S}} = O_P(\log pn^{-\tau}), \quad (2)$$

where  $\|\cdot\|_{\mathcal{S}}$  denotes the Hilbert–Schmidt norm, and the parameter  $\tau \in (0, 1/2]$  reflects the average sampling frequency, with larger values yielding denser observational points. Fang et al. (2024) developed the functional covariance estimation with theoretical guarantees by assuming generalized sub-Gaussian-type concentration inequalities for local linear smoothers  $\widehat{\Sigma}_{jk}(\cdot, \cdot)$ 's, resulting in an improved elementwise maximum rate:

$$\max_{1 \leq j, k \leq p} \|\widehat{\Sigma}_{jk} - \Sigma_{jk}\|_{\mathcal{S}} = O_P\{(\log p)^{1/2}n^{-\tau} + h^2\}, \quad (3)$$

where  $h > 0$  is the bandwidth parameter. However, it remains of theoretical interest to ask:

- What are the exact forms of such rates as functions of  $n$ ,  $p$ ,  $T_{ij}$ 's, and associated bandwidth parameters under cases with different sampling frequencies?
- Are these rates well-established in the sense of specifying the largest values of  $\tau$  and, compared to Zhang and Wang (2016), exhibiting any corresponding phase transition phenomena in the high-dimensional setting?

This paper aims to fill crucial theoretical gaps related to local linear smoothers frequently adopted in existing literature. Specifically, we present a systematic and unified non-asymptotic analysis of local linear smoothers for the mean and covariance functions to accommodate both sparsely and densely observed functional data in high dimensions. While our focus is not to introduce new methodologies for handling high-dimensional partially observed functional data, we make three new contributions as follows.

- First, we develop generalized sub-Gaussian-type concentration inequalities for each functional element of the mean and covariance estimators in both  $L_2$  norm and supremum norm. Compared to the asymptotic results in Zhang and Wang (2016), our non-asymptotic error bounds lead to the same rates of  $L_2$  convergence and uniform convergence, and reveal the same phase transition phenomena depending on the relative order of the average sampling frequency per subject to  $n^{1/4}$  for dense functional data. See Remarks 4 and 5.
- Second, we derive elementwise maximum rates of both  $L_2$  and uniform convergence for the mean and covariance estimators. Notably, we fundamentally improve the rates (2) and (3) assumed in existing literature in the sense of precisely specifying the largest values of  $\tau$  under cases with different sampling frequencies. These established rates in Theorems 6 and 7 serve as a foundational tool to provide theoretical guarantees for a set of aforementioned sparse high-dimensional functional models in the existing literature when dealing with the practical scenario of partially observed functional data in high dimensions.
- Third, with the presence of additional  $\log p$  terms to account for the high-dimensional effect in our established elementwise maximum rates, the scaled phase transitions

for high-dimensional dense functional data occur based on the ratios of the average sampling frequency per subject to  $n^{1/4}(\log p)^{-1/4}$ . This leads to a further partition of dense functional data into categories of “semi-dense” and “ultra-dense”, depending on whether the parametric rate  $(\log p)^{1/2}n^{-1/2}$  can be attained or not. With suitable choices of optimal bandwidths, we also present the optimal elementwise maximum rates from sparse to semi-dense to ultra-dense functional data, which correspondingly extend the optimal rates in Zhang and Wang (2016) to the high-dimensional setting. See Remarks 8 and 9.

**Outline of the paper.** Section 2 presents the nonparametric smoothing approach to estimate the mean and covariance functions. In Section 3, we investigate the non-asymptotic properties of the proposed local linear smoothers and discuss the associated phase transition phenomena. In Section 4, we outline a couple of applications of the non-asymptotic theory for the local linear smoothers. The established theoretical results are validated through simulations in Section 5. All technical proofs are relegated to the appendix.

**Notation.** We summarize here some notation to be used throughout the paper. For a positive integer  $q$ , we write  $[q] = \{1, \dots, q\}$ . For  $x, y \in \mathbb{R}$ , we write  $x \vee y = \max(x, y)$  and  $x \wedge y = \min(x, y)$ . We use  $I(\cdot)$  to denote an indicator function. We use  $\mathbf{I}_p$  to denote the  $p \times p$  identity matrix. Let  $L_2(\mathcal{U})$  be a Hilbert space of square-integrable functions on a compact interval  $\mathcal{U}$  equipped with the inner product  $\langle f, g \rangle = \int f(u)g(u)du$  for  $f(\cdot), g(\cdot) \in L_2(\mathcal{U})$  and the induced  $L_2$  norm  $\|\cdot\|_2 = \langle \cdot, \cdot \rangle^{1/2}$ . For any bivariate function  $\Phi(\cdot, \cdot)$  in  $L_2(\mathcal{U} \times \mathcal{U})$ , we also use  $\Phi$  to denote the linear operator induced from the kernel function  $\Phi(\cdot, \cdot)$ , that is, for any  $f(\cdot) \in L_2(\mathcal{U})$ ,  $\Phi(f)(\cdot) = \int \Phi(\cdot, v)f(v)dv \in L_2(\mathcal{U})$ , and denote its Hilbert–Schmidt norm by  $\{\int \int \Phi(u, v)^2 dudv\}^{1/2}$ . For two positive sequences  $\{a_n\}$  and  $\{b_n\}$ , we write  $a_n \lesssim b_n$  or  $b_n \gtrsim a_n$  if there exist a positive constant  $c$  such that  $\limsup_{n \rightarrow \infty} a_n/b_n \leq c$ . We write  $a_n \asymp b_n$  if and only if  $a_n \lesssim b_n$  and  $b_n \lesssim a_n$  hold simultaneously.

## 2. Methodology

Let  $\mathbf{X}_i(\cdot) = \{X_{i1}(\cdot), \dots, X_{ip}(\cdot)\}^T$  for  $i \in [n]$  be independently and identically distributed copies of  $\mathbf{X}(\cdot)$  defined on  $\mathcal{U}$  with mean  $\boldsymbol{\mu}(\cdot)$  and covariance  $\boldsymbol{\Sigma}(\cdot, \cdot)$ . For any  $i \in [n]$  and  $j \in [p]$ ,  $X_{ij}(\cdot)$  is not directly observable in practice. Instead it is observed, with random errors, at  $T_{ij}$  random time points,  $U_{ij1}, \dots, U_{ijT_{ij}} \in \mathcal{U}$ . Let  $Y_{ijt}$  be the observed value of  $X_{ij}(U_{ijt})$  satisfying

$$Y_{ijt} = X_{ij}(U_{ijt}) + \varepsilon_{ijt}, \tag{4}$$

where the errors  $\varepsilon_{ijt}$ 's, independent of  $X_{ij}$ 's, are independently and identically distributed copies of  $\varepsilon_j$  with  $\mathbb{E}(\varepsilon_j) = 0$  and  $\text{var}(\varepsilon_j) = \sigma_j^2 < \infty$ .

Based on the observed data  $\{(U_{ijt}, Y_{ijt}) : i \in [n], j \in [p], t \in [T_{ij}]\}$ , we present a unified procedure to estimate the mean functions  $\mu_j(\cdot)$ 's and the marginal- and cross-covariance functions  $\Sigma_{jk}(\cdot, \cdot)$ 's for both sparsely and densely observed functional data. In what follows, denote  $K_h(\cdot) = h^{-1}K(\cdot/h)$  for a univariate kernel  $K$  with bandwidth  $h > 0$ . For each  $j$ , a local linear smoother is firstly applied to  $\{(U_{ijt}, Y_{ijt}) : i \in [n], t \in [T_{ij}]\}$ , and hence the

estimated mean function is attained via  $\hat{\mu}_j(u) = \hat{b}_0$ , where

$$(\hat{b}_0, \hat{b}_1) = \arg \min_{b_0, b_1} \sum_{i=1}^n v_{ij} \sum_{t=1}^{T_{ij}} \left\{ Y_{ijt} - b_0 - b_1(U_{ijt} - u) \right\}^2 K_{h_{\mu, j}}(U_{ijt} - u).$$

The weight  $v_{ij}$  is attached to each observation for the  $i$ th subject and the  $j$ th functional variable such that  $\sum_{i=1}^n T_{ij} v_{ij} = 1$  (Zhang and Wang, 2016).

For each  $i \in [n]$ ,  $j, k \in [p]$ ,  $t \in [T_{ij}]$  and  $s \in [T_{ik}]$ , once the mean functions are estimated, let  $\Theta_{ij kts} = \{Y_{ijt} - \hat{\mu}_j(U_{ijt})\} \{Y_{iks} - \hat{\mu}_k(U_{iks})\}$  be the ‘‘raw covariance’’ between  $Y_{ijt}$  and  $Y_{iks}$ . Notice that  $\text{cov}(Y_{ijt}, Y_{iks}) = \Sigma_{jk}(U_{ijt}, U_{iks}) + \sigma_j^2 I(j = k) I(t = s)$ . To estimate the marginal-covariance function  $\Sigma_{jj}(\cdot, \cdot)$  for each  $j$  or the cross-covariance function  $\Sigma_{jk}(\cdot, \cdot)$  for each  $j \neq k$ , we employ local linear surface smoothers to the off-diagonals of the raw marginal-covariances  $(\Theta_{ij kts})_{1 \leq t \neq s \leq T_{ij}}$  or to the raw cross-covariances  $(\Theta_{ij kts})_{t \in [T_{ij}], s \in [T_{ik}]}$ . Specifically, we minimize

$$\sum_{i=1}^n w_{ijk} \sum_{(t,s) \in \mathcal{T}} \left\{ \Theta_{ij kts} - \beta_0 - \beta_1(U_{ijt} - u) - \beta_2(U_{iks} - v) \right\}^2 K_{h_{\Sigma, jk}}(U_{ijt} - u) K_{h_{\Sigma, jk}}(U_{iks} - v)$$

with respect to  $(\beta_0, \beta_1, \beta_2)$ , where the set  $\mathcal{T}$  equals to  $\{(t, s) : t \in [T_{ij}], s \in [T_{ij}], t \neq s\}$  if  $j = k$  or  $\{(t, s) : t \in [T_{ij}], s \in [T_{ik}]\}$  if  $j \neq k$ , and the weight  $w_{ijk}$  is assigned to each triplet  $(i, j, k)$  such that  $\sum_{i=1}^n T_{ij} \{T_{ik} - I(j = k)\} w_{ijk} = 1$ . See the weights to estimate marginal-covariance functions in Zhang and Wang (2016). The resulting marginal- or cross-covariance estimator is  $\hat{\Sigma}_{jk}(u, v) = \hat{\beta}_0$ . For ease of presentation, we assume that the mean functions  $\mu_j(\cdot)$ ’s are known in advance when discussing the concentration and convergence results related to the covariance estimators  $\hat{\Sigma}_{jk}(u, v)$ ’s in Section 3 below. However, it is noteworthy that all our discussions remain valid even when  $\mu_j(\cdot)$ ’s are unknown as long as a few additional technical assumptions are imposed.

Our estimation procedure allows general weighting schemes for  $\{v_i\}_{i \in [n]}$ ,  $\{w_{ijk}\}_{i \in [n], j, k \in [p]}$  such that two types of frequently-used schemes in existing literature are special cases of them. One type assigns the same weights to each observation (Yao et al., 2005a) with  $v_{ij} = (\sum_{i=1}^n T_{ij})^{-1}$  and  $w_{ijk} = [\sum_{i=1}^n T_{ij} \{T_{ik} - I(j = k)\}]^{-1}$ , so a subject with a larger number of observations receives more weights in total. The other type assigns the same weights to each subject (Li and Hsing, 2010), thus leading to  $v_{ij} = (nT_{ij})^{-1}$  and  $w_{ijk} = [nT_{ij} \{T_{ik} - I(j = k)\}]^{-1}$ .

**Remark 1** (i) Suppose that the estimated mean and covariance functions are evaluated at a grid of  $R \times R$  locations over  $\mathcal{U}^2$ . Under high-dimensional settings, it is apparent that our nonparametric smoothing approach suffers from high computational cost in kernel evaluations, particularly when estimating  $p(p+1)/2$  marginal- and cross-covariance functions. In a common practical scenario, where each  $X_{ij}(\cdot)$  is observed at the same set of time points  $U_1, \dots, U_{T_i} \in \mathcal{U}$  across  $j \in [p]$ , model (4) simplifies to

$$Y_{ijt} = X_{ij}(U_{it}) + \varepsilon_{ijt}, \quad t = 1, \dots, T_i,$$

which reduces the number of kernel evaluations from  $O(\sum_{i=1}^n \sum_{j=1}^p T_{ij} R)$  to  $O(\sum_{i=1}^n T_i R)$ , substantially accelerating the computation in high-dimensional settings. To further speed up

the computation, we can use the linear binning technique (Fan and Marron, 1994) to approximate the mean and covariance estimation. This would largely reduce the number of kernel evaluations to  $O(R)$ , and the number of operations (i.e. additions and multiplications) to  $O(np^2R^2 + p^2R^4 + p \sum_{i=1}^n T_i)$ , from  $O(p^2R^2 \sum_{i=1}^n T_i^2)$ . See the detailed implementation of binning and the associated computational complexity analysis in Fang et al. (2024). Our conducted numerical experiments show that such binned implementation offers significantly improved computational efficiency without sacrificing any estimation accuracy.

(ii) In a general scenario when  $X_{ij}(\cdot)$ 's are observed at different sets of time points, parallel estimation for  $j, k \in [p]$  can be employed, resulting in a more efficient procedure. In contrast, under certain lower-dimensional structural assumptions, the possible development of a nonparametric smoothing method for the joint estimation of components in  $\Sigma(\cdot, \cdot)$  becomes challenging in this general scenario, and is thus left for future research.

(iii) Due to the infinite-dimensional nature of functional data, it is standard practice to employ FPCA as a first dimension reduction step before subsequent modeling in the commonly-adopted multi-step estimation when dealing with high-dimensional functional models. See the detailed discussion of this application in Section 4.1. Specifically, implementing FPCA necessitates the nonparametric estimation of only marginal-covariance functions  $\Sigma_{jj}(\cdot, \cdot)$  across  $j \in [p]$ , which can be easily paralleled for fast computation.

### 3. Theory

Before presenting the concentration and convergence results, we impose the following regularity assumptions.

**Assumption 1** For each  $i \in [n]$  and  $j \in [p]$ ,  $X_{ij}(\cdot)$  is a sub-Gaussian random process and  $\varepsilon_{ij}$  is a sub-Gaussian random variable, that is, there exists some positive constant  $c$  such that  $\mathbb{E}\{\exp(\langle x, X_{ij} - \mu_j \rangle)\} \leq \exp\{2^{-1}c^2\langle x, \Sigma_{jj}(x) \rangle\}$  for all  $x(\cdot) \in L^2(\mathcal{U})$  and  $\mathbb{E}\{\exp(\varepsilon_{ij}z)\} \leq \exp(c^2\sigma_j^2z^2/2)$  for all  $z \in \mathbb{R}$ .

**Assumption 2** For each  $i \in [n]$  and  $j \in [p]$ , under the sparse design,  $T_{ij} \leq T_0 < \infty$ , and, under the dense design,  $T_{ij} \rightarrow \infty$  and there exists some positive constant  $\bar{c}$  such that  $\max_{i,j} T_{ij} (\min_{i,j} T_{ij})^{-1} \leq \bar{c}$ .

**Assumption 3** Under the dense design, there exists some positive constant  $c_0$  such that  $\max_{i,j} v_{ij} (\min_{i,j} v_{ij})^{-1} \leq c_0$  and  $\max_{i,j,k} w_{ijk} (\min_{i,j,k} w_{ijk})^{-1} \leq c_0$ .

**Assumption 4** (i) Let  $\{U_{ijt} : i \in [n], j \in [p], t \in [T_{ij}]\}$  be independently and identically distributed copies of a random variable  $U$  defined on  $\mathcal{U}$ . The density  $f_U(\cdot)$  of  $U$  satisfies  $0 < m_f \leq \inf_{\mathcal{U}} f_U(u) \leq \sup_{\mathcal{U}} f_U(u) \leq M_f < \infty$  for some positive constants  $m_f$  and  $M_f$ ; (ii)  $\mathbf{X}(\cdot)$ ,  $U$  and  $\{\varepsilon_j\}_{j \in [p]}$  are mutually independent.

**Assumption 5** Let  $B_{jk} = [(k-1)\tilde{T}_j^{-1}, k\tilde{T}_j^{-1}]$  for  $k \in [\tilde{T}_j]$  with  $\tilde{T}_j = \max_i T_{ij}$ , there exists some constant  $C > 0$  such that the cardinality  $\#\{U_{ijt} : U_{ijt} \in B_{jk}, t \in [T_{ij}]\} \leq C$  for each  $i \in [n], j \in [p]$  and  $k \in [\tilde{T}_j]$ .

**Assumption 6** (i)  $K(\cdot)$  is a symmetric probability density function defined on  $[-1, 1]$  with  $\int u^2 K(u) du < \infty$  and  $\int K(u)^2 du < \infty$ . (ii)  $K(\cdot)$  is Lipschitz continuous: there exists some positive constant  $L$  such that  $|K(u) - K(v)| \leq L|u - v|$  for any  $u, v \in [-1, 1]$ .

**Assumption 7** (i)  $\partial^2 \mu_j(u)/\partial u^2$  is uniformly bounded over  $u \in \mathcal{U}$  and  $j \in [p]$ ; (ii)  $\partial^2 \Sigma_{jk}(u, v)/\partial u^2$ ,  $\partial^2 \Sigma_{jk}(u, v)/\partial u \partial v$ , and  $\partial^2 \Sigma_{jk}(u, v)/\partial v^2$  are uniformly bounded over  $(u, v) \in \mathcal{U}^2$  and  $(j, k) \in [p]^2$ .

The sub-Gaussianities in Assumption 1 for both Hilbert space-valued random elements  $X_{ij}(\cdot)$ 's and random errors  $\varepsilon_{ij}$ 's together imply that the observations  $Y_{ijt}$ 's in (4) are sub-Gaussian, which plays a crucial role in deriving our subsequent concentration inequalities. The dense case in Assumption 2 corresponds to a common practical scenario, where the sampling frequencies  $T_{ij}$ 's are of the same order across  $i \in [n]$  and  $j \in [p]$ . Under such a scenario, Assumption 3 is automatically satisfied by two frequently-used weighting schemes including ‘‘equal weight per observation’’ and ‘‘equal weight per subject’’. Assumption 5 means that all observational time points are distributed in the sense of ‘‘uniformly on  $\mathcal{U}$ ’’. This prevents the occurrence of an extreme case where a large number of time points are concentrated in some small areas while leaving too few points in other regions. Assumptions 4, 6 and 7 are standard in the literature of local linear smoothing for functional data (Yao et al., 2005a; Zhang and Wang, 2016) adaptable to the multivariate setting.

**Theorem 2** Suppose that Assumptions 1-6 hold. For each  $j \in [p]$ , let  $\gamma_{n,T,h,j} = n(1 \wedge \bar{T}_{\mu,j} h_{\mu,j})$  with the corresponding average sampling frequency per subject  $\bar{T}_{\mu,j} = n^{-1} \sum_{i=1}^n T_{ij}$ , then there exist some positive constants  $c_1, c_2$  (independent of  $n, p, \bar{T}_{\mu,j}$ 's) and arbitrarily small  $\epsilon_1 > 0$  such that for any  $\delta \in (0, 1]$ ,

$$\mathbb{P}(\|\hat{\mu}_j - \tilde{\mu}_j\|_2 \geq \delta) \leq c_2 \exp(-c_1 \gamma_{n,T,h,j} \delta^2), \quad (5)$$

$$\mathbb{P}\left\{\sup_{u \in \mathcal{U}} |\hat{\mu}_j(u) - \tilde{\mu}_j(u)| \geq \delta\right\} \leq \frac{c_2 (n^{\epsilon_1} \gamma_{n,T,h,j})^{1/2}}{h_{\mu,j}^2} \exp(-c_1 \gamma_{n,T,h,j} \delta^2), \quad (6)$$

where  $\tilde{\mu}_j(u)$  is a deterministic univariate function that converges to  $\mu_j(u)$  as  $h_{\mu,j} \rightarrow 0$ . See (A.1) in Appendix A for the exact form of  $\tilde{\mu}_j(u)$ .

**Theorem 3** Suppose that Assumptions 1-6 hold. For each  $j, k \in [p]$ , let  $\nu_{n,T,h,jk} = n(1 \wedge \bar{T}_{\Sigma,jk}^2 h_{\Sigma,jk}^2)$  with the corresponding average sampling frequency per subject being  $\bar{T}_{\Sigma,jk} = [n^{-1} \sum_{i=1}^n T_{ij} \{T_{ik} - I(j=k)\}]^{1/2}$ , then there exist some positive constants  $c_3, c_4$  (independent of  $n, p, \bar{T}_{\Sigma,jk}$ 's) and arbitrarily small  $\epsilon_2 > 0$  such that for any  $\delta \in (0, 1]$ ,

$$\mathbb{P}\left(\|\widehat{\Sigma}_{jk} - \tilde{\Sigma}_{jk}\|_S \geq \delta\right) \leq c_4 \exp(-c_3 \nu_{n,T,h,jk} \delta^2), \quad (7)$$

$$\mathbb{P}\left\{\sup_{(u,v) \in \mathcal{U}^2} |\widehat{\Sigma}_{jk}(u,v) - \tilde{\Sigma}_{jk}(u,v)| \geq \delta\right\} \leq \frac{c_4 n^{\epsilon_2} \nu_{n,T,h,jk}}{h_{\Sigma,jk}^6} \exp(-c_3 \nu_{n,T,h,jk} \delta^2), \quad (8)$$

where  $\tilde{\Sigma}_{jk}(u, v)$  is a deterministic bivariate function that converges to  $\Sigma_{jk}(u, v)$  as  $h_{\Sigma,jk} \rightarrow 0$ . See (A.19) in Appendix A for the exact form of  $\tilde{\Sigma}_{jk}(u, v)$ .

**Remark 4** *The concentration inequalities in Theorems 2 and 3 imply that  $\hat{\mu}_j$  and  $\hat{\Sigma}_{jk}$  are nicely concentrated around  $\tilde{\mu}_j$  and  $\tilde{\Sigma}_{jk}$ , respectively, in both  $L_2$  norm and supremum norm with generalized sub-Gaussian-type tail behaviors. It is worth mentioning that such  $L_2$  and uniform concentration results are derived based on the local concentration inequalities of  $\hat{\mu}_j(u)$  and  $\hat{\Sigma}_{j,k}(u, v)$  for fixed interior points  $u, v \in \mathcal{U}$ , which enjoy the same tail behaviors as (5) and (7). Besides being fundamental to derive elementwise maximum error bounds that are essential for further convergence analysis under high-dimensional settings, these non-asymptotic results lead to the same rates of  $L_2$  convergence and uniform convergence compared to those in Zhang and Wang (2016). Specifically, under extra Assumption 7, it holds that*

$$\begin{aligned} \|\hat{\mu}_j - \mu_j\|_2 &= O_P\{n^{-1/2} + (n\bar{T}_{\mu,j}h_{\mu,j})^{-1/2} + h_{\mu,j}^2\}, \\ \sup_{u \in \mathcal{U}} |\hat{\mu}_j(u) - \mu_j(u)| &= O_P[(\log n)^{1/2}n^{-1/2}\{1 + (\bar{T}_{\mu,j}h_{\mu,j})^{-1/2}\} + h_{\mu,j}^2], \\ \|\hat{\Sigma}_{jk} - \Sigma_{jk}\|_S &= O_P\{n^{-1/2} + (n\bar{T}_{\Sigma,jk}^2h_{\Sigma,jk}^2)^{-1/2} + h_{\Sigma,jk}^2\}, \\ \sup_{(u,v) \in \mathcal{U}^2} |\hat{\Sigma}_{jk}(u, v) - \Sigma_{jk}(u, v)| &= O_P[(\log n)^{1/2}n^{-1/2}\{1 + (\bar{T}_{\Sigma,jk}^2h_{\Sigma,jk}^2)^{-1/2}\} + h_{\Sigma,jk}^2]. \end{aligned}$$

**Remark 5** *The above rates of convergence reveal interesting phase transition phenomena depending on the ratio of the average sampling frequency per subject  $\bar{T}_{\mu,j}$  (or  $\bar{T}_{\Sigma,jk}$ ) to  $n^{1/4}$ . In the following, we use different rates of  $L_2$  convergence for  $\hat{\mu}_j$  and  $\hat{\Sigma}_{jk}$  to illustrate a systematic partition of partially observed functional data into three categories:*

1. Under the sparse design, when  $h_{\mu,j} \asymp n^{-1/5}$ ,

$$\|\hat{\mu}_j - \mu_j\|_2 = O_P(n^{-1/2}h_{\mu,j}^{-1/2} + h_{\mu,j}^2) = O_P(n^{-2/5});$$

when  $h_{\Sigma,jk} \asymp n^{-1/6}$ ,

$$\|\hat{\Sigma}_{jk} - \Sigma_{jk}\|_S = O_P(n^{-1/2}h_{\Sigma,jk}^{-1} + h_{\Sigma,jk}^2) = O_P(n^{-1/3}).$$

2. Under the dense design, when  $\bar{T}_{\mu,j}n^{-1/4} \rightarrow 0$  with  $h_{\mu,j} \asymp (n\bar{T}_{\mu,j})^{-1/5}$ ,

$$\|\hat{\mu}_j - \mu_j\|_2 = O_P(n^{-1/2}\bar{T}_{\mu,j}^{-1/2}h_{\mu,j}^{-1/2} + h_{\mu,j}^2) = O_P\{(n\bar{T}_{\mu,j})^{-2/5}\};$$

when  $\bar{T}_{\Sigma,jk}n^{-1/4} \rightarrow 0$  with  $h_{\Sigma,jk} \asymp (n\bar{T}_{\Sigma,jk}^2)^{-1/6}$ ,

$$\|\hat{\Sigma}_{jk} - \Sigma_{jk}\|_S = O_P(n^{-1/2}\bar{T}_{\Sigma,jk}^{-1}h_{\Sigma,jk}^{-1} + h_{\Sigma,jk}^2) = O_P\{(n\bar{T}_{\Sigma,jk}^2)^{-1/3}\}.$$

3. Under the dense design, when  $\bar{T}_{\mu,j}n^{-1/4} \rightarrow \tilde{c}$  (some positive constant) with  $h_{\mu,j} \asymp n^{-1/4}$  or  $\bar{T}_{\mu,j}n^{-1/4} \rightarrow \infty$  with  $h_{\mu,j} = o(n^{-1/4})$  and  $\bar{T}_{\mu,j}h_{\mu,j} \rightarrow \infty$ ,

$$\|\hat{\mu}_j - \mu_j\|_2 = O_P(n^{-1/2});$$

when  $\bar{T}_{\Sigma,jk}n^{-1/4} \rightarrow \tilde{c}$  with  $h_{\Sigma,jk} \asymp n^{-1/4}$  or  $\bar{T}_{\Sigma,jk}n^{-1/4} \rightarrow \infty$  with  $h_{\Sigma,jk} = o(n^{-1/4})$  and  $\bar{T}_{\Sigma,jk}h_{\Sigma,jk} \rightarrow \infty$ ,

$$\|\hat{\Sigma}_{jk} - \Sigma_{jk}\|_S = O_P(n^{-1/2}).$$

As  $\bar{T}_{\mu,j}$  and  $\bar{T}_{\Sigma,jk}$  grow very fast, case 3 results in the root- $n$  rate complying with the parametric rate for fully observed functional data. As  $\bar{T}_{\mu,j}$  and  $\bar{T}_{\Sigma,jk}$  grow moderately fast, case 2 corresponds to the optimal minimax rates (Zhang and Wang, 2016), which are slower than root- $n$  but faster than the counterparts for sparsely observed functional data. Our established convergence rates in cases 1, 2 and 3 allow free choices of  $(j, k)$ , and are respectively consistent to those of the mean and covariance estimators under categories of “sparse”, “semi-dense” and “ultra-dense” univariate functional data introduced in Zhang and Wang (2016).

**Theorem 6** Suppose that the assumptions in Theorem 2 and Assumption 7(i) hold, and  $(\min_j \gamma_{n,T,h,j})^{-1} \log p \rightarrow 0$ ,  $\max_j h_{\mu,j} \rightarrow 0$  as  $n, p \rightarrow \infty$ . It then holds that

$$\max_{j \in [p]} \|\hat{\mu}_j - \mu_j\|_2 = O_P \left\{ \left( \frac{\log p}{\min_j \gamma_{n,T,h,j}} \right)^{1/2} + \max_j h_{\mu,j}^2 \right\}, \quad (9)$$

and, if  $\min_j h_{\mu,j} \asymp \{\log(p \vee n)/n\}^{\kappa_1}$  for some  $\kappa_1 \in (0, 1/2]$ ,

$$\max_{j \in [p]} \sup_{u \in \mathcal{U}} |\hat{\mu}_j(u) - \mu_j(u)| = O_P \left[ \left\{ \frac{\log(p \vee n)}{\min_j \gamma_{n,T,h,j}} \right\}^{1/2} + \max_j h_{\mu,j}^2 \right]. \quad (10)$$

**Theorem 7** Suppose that the assumptions in Theorem 3 and Assumption 7(ii) hold, and  $(\min_{j,k} \nu_{n,T,h,jk})^{-1} \log p \rightarrow 0$ ,  $\max_{j,k} h_{\Sigma,jk} \rightarrow 0$  as  $n, p \rightarrow \infty$ . It then holds that

$$\max_{j,k \in [p]} \|\hat{\Sigma}_{jk} - \Sigma_{jk}\|_{\mathcal{S}} = O_P \left\{ \left( \frac{\log p}{\min_{j,k} \nu_{n,T,h,jk}} \right)^{1/2} + \max_{j,k} h_{\Sigma,jk}^2 \right\}, \quad (11)$$

and, if  $\min_{j,k} h_{\Sigma,jk} \asymp \{\log(p \vee n)/n\}^{\kappa_2}$  for some  $\kappa_2 \in (0, 1/2]$ ,

$$\max_{j,k \in [p]} \sup_{(u,v) \in \mathcal{U}^2} |\hat{\Sigma}_{jk}(u,v) - \Sigma_{jk}(u,v)| = O_P \left[ \left\{ \frac{\log(p \vee n)}{\min_{j,k} \nu_{n,T,h,jk}} \right\}^{1/2} + \max_{j,k} h_{\Sigma,jk}^2 \right]. \quad (12)$$

We observe that the elementwise maximum rates of  $L_2$  convergence and uniform convergence are governed by both dimensionality parameters  $(n, p, \{\bar{T}_{\mu,j}\}_{j \in [p]}, \{\bar{T}_{\Sigma,jk}\}_{j,k \in [p]})$  and internal parameters  $(\{h_{\mu,j}\}_{j \in [p]}, \{h_{\Sigma,jk}\}_{j,k \in [p]})$ . Each convergence rate is composed of two terms reflecting our familiar variance-bias tradeoff in nonparametric statistics. It is easy to see that the variance terms are determined by the least frequently sampled and smoothed components, that is the smallest  $\bar{T}_{\mu,j}$  (or  $\bar{T}_{\Sigma,jk}$ ) and  $h_{\mu,j}$  (or  $h_{\Sigma,jk}$ ) across  $j, k$ , whereas the highest level of smoothness with the largest  $h_{\mu,j}$  (or  $h_{\Sigma,jk}$ ) controls the bias terms.

**Remark 8** To facilitate further discussion, we consider the simplified setting where  $\bar{T}_{\mu,j} \asymp \bar{T}_{\mu}$ ,  $h_{\mu,j} \asymp h_{\mu}$  and  $\bar{T}_{\Sigma,jk} \asymp \bar{T}_{\Sigma}$ ,  $h_{\Sigma,jk} \asymp h_{\Sigma}$  for each  $j, k$ . Compared to cases 1–3 above, the corresponding elementwise maximum rates of convergence for  $\{\hat{\mu}_j\}_{j \in [p]}$  (or  $\{\hat{\Sigma}_{jk}\}_{j,k \in [p]}$ ) in Theorem 6 (or Theorem 7) reveal scaled phase transitions for dense functional data depending on the relative order of  $\bar{T}_{\mu}$  (or  $\bar{T}_{\Sigma}$ ) to  $n^{1/4}(\log p)^{-1/4}$  instead of  $n^{1/4}$ . In the following, we use elementwise maximum rates of  $L_2$  convergence to illustrate the phase transition phenomena and the optimal estimation from sparse to dense functional data in high dimensions. In terms of uniform convergence, the same phenomena occur as long as  $p \gtrsim n$ .

(i) Under the sparse design, when  $h_\mu \asymp (\log p)^{1/5} n^{-1/5}$ ,

$$\max_j \|\hat{\mu}_j - \mu_j\|_2 = O_P \left\{ \left( \frac{\log p}{nh_\mu} \right)^{1/2} + h_\mu^2 \right\} = O_P \left\{ \left( \frac{\log p}{n} \right)^{2/5} \right\};$$

when  $h_\Sigma \asymp (\log p)^{1/6} n^{-1/6}$ ,

$$\max_{j,k} \|\hat{\Sigma}_{jk} - \Sigma_{jk}\|_S = O_P \left\{ \left( \frac{\log p}{nh_\Sigma^2} \right)^{1/2} + h_\Sigma^2 \right\} = O_P \left\{ \left( \frac{\log p}{n} \right)^{1/3} \right\}.$$

(ii) Under the dense design, when  $\bar{T}_\mu (\log p)^{1/4} n^{-1/4} \rightarrow 0$  with  $h_\mu \asymp (\log p)^{1/5} (n\bar{T}_\mu)^{-1/5}$ ,

$$\max_j \|\hat{\mu}_j - \mu_j\|_2 = O_P \left\{ \left( \frac{\log p}{n\bar{T}_\mu h_\mu} \right)^{1/2} + h_\mu^2 \right\} = O_P \left\{ \left( \frac{\log p}{n\bar{T}_\mu} \right)^{2/5} \right\}; \quad (13)$$

when  $\bar{T}_\Sigma (\log p)^{1/4} n^{-1/4} \rightarrow 0$  with  $h_\Sigma \asymp (\log p)^{1/6} (n\bar{T}_\Sigma^2)^{-1/6}$ ,

$$\max_{j,k} \|\hat{\Sigma}_{jk} - \Sigma_{jk}\|_S = O_P \left\{ \left( \frac{\log p}{n\bar{T}_\Sigma^2 h_\Sigma^2} \right)^{1/2} + h_\Sigma^2 \right\} = O_P \left\{ \left( \frac{\log p}{n\bar{T}_\Sigma^2} \right)^{1/3} \right\}. \quad (14)$$

(iii) Under the dense design, when  $\bar{T}_\mu (\log p)^{1/4} n^{-1/4} \rightarrow \tilde{c}$  with  $h_\mu \asymp (\log p)^{1/4} n^{-1/4}$  or  $\bar{T}_\mu (\log p)^{1/4} n^{-1/4} \rightarrow \infty$  with  $h_\mu = o\{(\log p)^{1/4} n^{-1/4}\}$  and  $\bar{T}_\mu h_\mu \rightarrow \infty$ ,

$$\max_j \|\hat{\mu}_j - \mu_j\|_2 = O_P \left\{ \left( \frac{\log p}{n} \right)^{1/2} \right\};$$

when  $\bar{T}_\Sigma (\log p)^{1/4} n^{-1/4} \rightarrow \tilde{c}$  with  $h_\Sigma \asymp (\log p)^{1/4} n^{-1/4}$  or  $\bar{T}_\Sigma (\log p)^{1/4} n^{-1/4} \rightarrow \infty$  with  $h_\Sigma = o\{(\log p)^{1/4} n^{-1/4}\}$  and  $\bar{T}_\Sigma h_\Sigma \rightarrow \infty$ ,

$$\max_{j,k} \|\hat{\Sigma}_{jk} - \Sigma_{jk}\|_S = O_P \left\{ \left( \frac{\log p}{n} \right)^{1/2} \right\}.$$

**Remark 9** In a similar spirit to the partitioned three categories for univariate functional data (see cases 1, 2 and 3 above), we can also term the high-dimensional partially observed functional data in cases (i), (ii), and (iii) as “sparse”, “semi-dense”, and “ultra-dense”, respectively. The main difference lies in the presence of additional  $\log p$  terms to account for the high-dimensional effect.

- As  $\bar{T}_\mu$  and  $\bar{T}_\Sigma$  grow at least in the order of  $n^{1/4} (\log p)^{-1/4}$ , the attained optimal rate  $(\log p)^{1/2} n^{-1/2}$  is identical to that for the fully observed functional data (Zapata et al., 2022), presenting that the theory for high-dimensional ultra-dense functional data falls in the parametric paradigm.
- As  $\bar{T}_\mu$  and  $\bar{T}_\Sigma$  diverge slower than  $n^{1/4} (\log p)^{-1/4}$ , if we let  $h_\mu \asymp (\log p)^{1/5} (n\bar{T}_\mu)^{-1/5}$  and  $h_\Sigma \asymp (\log p)^{1/6} (n\bar{T}_\Sigma^2)^{-1/6}$  to balance the corresponding variance and bias terms, the optimal rates for high-dimensional semi-dense functional data are respectively achieved in (13) and (14). These rates degenerate to the minimax rates in case 2 when  $p$

is fixed. With the choice of elementwise optimal bandwidths  $h_\mu \asymp (n\bar{T}_\mu)^{-1/5}$  and  $h_\Sigma \asymp (n\bar{T}_\Sigma^2)^{-1/6}$ , we obtain

$$\max_j \|\hat{\mu}_j - \mu_j\|_2 = O_P\{(\log p)^{1/2}(n\bar{T}_\mu)^{-2/5}\}, \quad \max_{j,k} \|\hat{\Sigma}_{jk} - \Sigma_{jk}\| = O_P\{(\log p)^{1/2}(n\bar{T}_\Sigma)^{-1/3}\},$$

which are respectively slower than the optimal rates in (13) and (14). Such discussion applies analogously to the sparse functional setting. See cases 1 and (i).

- Compared to the asymptotic results for cases 1, 2 and 3 under a fixed  $p$  scenario, the high-dimensionality in cases (i), (ii) and (iii) leads to the scaled phase transitions, optimal selected bandwidths, and corresponding optimal rates, each of which is up to a factor of  $\log p$  at some polynomial order.

## 4. Applications

In this section, we outline three applications of our established non-asymptotic results for the local linear smoothers under high dimensional settings.

### 4.1 Estimation Under the FPCA Framework

A standard procedure towards the estimation of models involving high-dimensional functional data consists of two or three steps. Due to the infinite-dimensionality of functional data, the first step performs dimension reduction via, e.g., FPCA, to approximate each  $X_{ij}(\cdot)$  by the  $d_j$ -dimensional truncation. This effectively transforms the problem of modeling the  $p$ -vector of functional variables into that of modeling the  $(\sum_{j=1}^p d_j)$ -vector of FPC scores. To overcome the difficulties caused by high-dimensionality, some functional sparsity assumptions are commonly imposed, which results in the estimation under block sparsity constraints in the second step. Examples include the group-lasso penalized least squares estimation in regression setups (Fan et al., 2015; Kong et al., 2016; Wang et al., 2022), the group graphical lasso in functional graphical model estimation (Qiao et al., 2019; Solea and Li, 2022; Zapata et al., 2022) and other related applications mentioned in Section 1. Finally, the third step re-transforms block sparse estimates obtained in the second step to functional sparse estimates via estimated principal component functions obtained in the first step. In functional graphical model estimation, the third step is no longer required. Building upon established theoretical results, this section presents some non-asymptotic results within the FPCA framework, which are crucial not only in their own right but also in providing the theoretical support for such multi-step estimation procedure.

For each  $j \in [p]$ , the standard dimension reduction method performs Karhunen-Loève expansion of each target trajectory  $X_{ij}(\cdot)$  and truncates the expansion to the first  $d_j$  terms, which serves as the foundation of FPCA:

$$X_{ij}(\cdot) = \mu_j(\cdot) + \sum_{l=1}^{\infty} \xi_{ijl} \phi_{jl}(\cdot) \approx \mu_j(\cdot) + \sum_{l=1}^{d_j} \xi_{ijl} \phi_{jl}(\cdot), \quad (15)$$

where the coefficients  $\xi_{ijl} = \langle X_{tj} - \mu_j, \phi_{jl} \rangle$  for  $l \geq 1$ , namely FPC scores, correspond to a sequence of random variables with  $\mathbb{E}(\xi_{ijl}) = \lambda_{jl}$  and  $\text{cov}(\xi_{ijl}, \xi_{ijl'}) = \lambda_{jl} I(l \neq l')$

and  $\lambda_{j1} \geq \lambda_{j2} \geq \dots > 0$  are the eigenvalues of  $\Sigma_{jj}(u, v)$  and  $\phi_{j1}(\cdot), \phi_{j2}(\cdot), \dots$  are the corresponding eigenfunctions. To implement FPCA based on partially observed functional data, we carry out an eigenanalysis of the local linear smoothers  $\hat{\Sigma}_{jj}(u, v)$  that leads to estimated eigenvalue/eigenfunction pairs  $(\hat{\lambda}_{jl}, \hat{\phi}_{jl}(\cdot))$  for  $j \in [p]$  and  $l \in [d_j]$ . Based on Theorem 7, we can establish the elementwise maximum rates for estimated eigenvalues and eigenfunctions in the following proposition.

**Proposition 10** *Suppose that the assumptions in Theorem 7 hold and  $\lambda_{j1} > \lambda_{j2} > \dots > 0$  for each  $j \in [p]$ . Let  $\delta_{jl} = \min_{k \in [l]} \{\lambda_{jk} - \lambda_{j(k+1)}\}$  for  $j \in [p]$  and  $l \in [d_j]$ . Then we have*

$$\max_{j \in [p], l \in [d_j]} \left\{ |\hat{\lambda}_{jl} - \lambda_{jl}| + \delta_{jl} \|\hat{\phi}_{jl} - \phi_{jl}\|_2 \right\} = O_P \left\{ \left( \frac{\log p}{\min_{j,k} \nu_{n,T,h,jk}} \right)^{1/2} + \max_{j,k} h_{\Sigma,jk}^2 \right\}.$$

Proposition 10 can be used to provide the theoretical guarantees for the first and third steps under high-dimensional settings. In the second step, the main target is to implement the block regularized estimation based on the estimated FPC scores. Under the dense design with  $T_{ij} \rightarrow \infty$ , the estimated FPC scores  $\langle X_{ij} - \hat{\mu}_j, \hat{\phi}_{jl} \rangle$  can be well approximated by the numerical integration based on  $\{U_{ijt}, Y_{ijt}, \hat{\phi}_{jl}(U_{ijt})\}$  for  $t \in [T_{ij}]$ . To be specific, we can employ a Trapezoid rule-based numerical integration to estimate FPC scores

$$\hat{\xi}_{ijl}^{(1)} = \sum_{t=2}^{T_{ij}} \frac{\{Y_{ij(t-1)} - \hat{\mu}_j(U_{ij(t-1)})\} \hat{\phi}_{jl}(U_{ij(t-1)}) + \{Y_{ijt} - \hat{\mu}_j(U_{ijt})\} \hat{\phi}_{jl}(U_{ijt})}{2} |U_{ijt} - U_{ij(t-1)}|. \quad (16)$$

However, such numerical integration approach fails under the sparse design with  $T_{ij} \leq T_0 < \infty$ . We instead employ the principal components analysis through conditional expectation (PACE) method (Yao et al., 2005a) to estimate FPC scores. For each  $j \in [p]$ , under the assumption that  $\xi_{ijl}$  and  $\varepsilon_{ijt}$  in (4) are jointly Gaussian, the PACE estimation of the FPC scores for the  $i$ th subject given the data from the individual reduces to the estimated conditional expectation

$$\hat{\xi}_{ijl}^{(2)} = \hat{\mathbb{E}}(\xi_{ijl} | \tilde{\mathbf{Y}}_{ij}) = \hat{\lambda}_{jl} \hat{\phi}_{ijl}^T \hat{\Sigma}_{\mathbf{Y}_{ij}}^{-1} (\tilde{\mathbf{Y}}_{ij} - \hat{\boldsymbol{\mu}}_{ij}), \quad (17)$$

where we write  $\tilde{\mathbf{Y}}_{ij} = (Y_{ij1}, \dots, Y_{ijT_{ij}})^T$ ,  $\hat{\Sigma}_{\mathbf{Y}_{ij}}$  is a  $T_{ij} \times T_{ij}$  matrix with its  $(t, t')$ -th entry  $(\hat{\Sigma}_{\mathbf{Y}_{ij}})_{t,t'} = \hat{\Sigma}_{jj}(U_{ijt}, U_{ijt'}) + \hat{\sigma}_j^2 I(t = t')$ ,  $\hat{\phi}_{ijl} = \{\hat{\phi}_{jl}(U_{ij1}), \dots, \hat{\phi}_{jl}(U_{ijT_{ij}})\}^T$ , and  $\hat{\boldsymbol{\mu}}_{ij} = (\hat{\mu}_j(U_{ij1}), \dots, \hat{\mu}_j(U_{ijT_{ij}}))^T$ . See Yao et al. (2005a) for details on the estimate  $\hat{\sigma}_j^2$  of  $\sigma_j$ .

For each  $j, k \in [p]$ ,  $l \in [d_j]$  and  $m \in [d_k]$ , let  $\sigma_{jklm}^{(h)} = \mathbb{E}(\xi_{ijl}^{(h)} \xi_{ikm}^{(h)})$  and its sample estimator be  $\hat{\sigma}_{jklm}^{(h)} = n^{-1} \sum_{i=1}^n \hat{\xi}_{ijl}^{(h)} \hat{\xi}_{ikm}^{(h)}$  for  $h = 1$  (dense case) and 2 (sparse case). To theoretically support the second step, it is essential to establish the elementwise maximum rates for  $\{\hat{\sigma}_{jklm}^{(h)}\}$  with  $h = 1, 2$ , i.e. the convergence rate of

$$\max_{j,k \in [p], l \in [d_j], m \in [d_k]} \frac{|\hat{\sigma}_{jklm}^{(h)} - \sigma_{jklm}^{(h)}|}{\Lambda_{jklm}^{(h)}},$$

where  $\Lambda_{jklm}^{(h)}$  represents some normalization term that may depend on  $\delta_{jl}$  and  $\delta_{km}$ . Our established elementwise maximum rate in Theorem 7 may still be applicable in this context.

However, this topic is beyond the scope of the current paper and is thus left for future research.

In addition, building upon the expansion in (15) and estimated FPC scores in (16) or (17), we can estimate the target trajectories for prediction or subsequent modeling

$$\widehat{X}_{ij}^{(h)}(\cdot) = \widehat{\mu}_j(\cdot) + \sum_{l=1}^{d_j} \widehat{\xi}_{ijl}^{(h)} \widehat{\phi}_{jl}(\cdot), \quad i \in [n], j \in [p], h = 1, 2. \quad (18)$$

We leave the development of non-asymptotic results for (18) as future research. Such results could possibly be useful for the application of nonparametric functional graphical model estimation (Li and Solea, 2018) when handling the practical partially observed functional scenario in high dimensions.

## 4.2 Sparse FPCA

While componentwise FPCA in Section 4.1 may fail to model the correlation between components in  $X_{i1}(\cdot), \dots, X_{ij}(\cdot)$ , more effective dimension reduction can be achieved by leveraging the correlation information across different components, such as multivariate FPCA (Happ and Greven, 2018) (for fixed  $p$ ) and sparse FPCA (Hu and Yao, 2022), which incorporates the notation of sparsity in multivariate statistics into the functional setting to accommodate high-dimensional functional data. The sparsity structure motivates a thresholding rule that is easy to compute by exploiting the relationship between the univariate orthonormal basis representation for infinite-dimensional processes and multivariate Karhunen-Loève expansion in the form of

$$\mathbf{X}_i(\cdot) = \boldsymbol{\mu}(\cdot) + \sum_{k=1}^{\infty} \zeta_{ik} \boldsymbol{\psi}_k(\cdot),$$

where  $\zeta_{ik} = \sum_{j=1}^p \langle X_{ij} - \mu_j, \boldsymbol{\psi}_{kj} \rangle$ ,  $\tilde{\lambda}_k$  and  $\boldsymbol{\psi}_k(\cdot) = \{\boldsymbol{\psi}_{k1}(\cdot), \dots, \boldsymbol{\psi}_{kp}(\cdot)\}^T$  for  $k = 1, \dots, \infty$  are the eigenvalues and the corresponding eigenfunctions of  $\boldsymbol{\Sigma}(u, v)$ , respectively, satisfying  $\int_{\mathcal{U}} \boldsymbol{\Sigma}(u, v) \boldsymbol{\psi}_k(v) dv = \tilde{\lambda}_k \boldsymbol{\psi}_k(u)$ . Specifically, provided with a complete and orthonormal basis  $\{b_l(\cdot) : l = 1, \dots, \infty\}$ , each random process is represented as  $X_{ij}(\cdot) = \mu_j(\cdot) + \sum_{l=1}^{\infty} \theta_{ijl} b_l(\cdot)$ , where the basis coefficients  $\theta_{ijl} = \langle X_{ij} - \mu_j, b_l \rangle$ . Denote  $\boldsymbol{\psi}_{kj}(\cdot) = \sum_{l=1}^{\infty} \eta_{kjl} b_l(\cdot)$ , where  $\eta_{kjl} = \langle \boldsymbol{\psi}_{kj}, b_l \rangle$ . According to Proposition 1 in Hu and Yao (2022), we have

$$\sum_{j'=1}^p \sum_{l'=1}^{\infty} \text{cov}(\theta_{ijl}, \theta_{ij'l'}) \eta_{kjl} = \tilde{\lambda}_k \eta_{kjl}, \quad i \in [n], j \in [p], k, l = 1, 2, \dots$$

Under the practical scenario (4) with  $T_{ij} \rightarrow \infty$  (dense case), Hu and Yao (2022) proposed to estimate  $\theta_{ijl}$  by  $\widehat{\theta}_{ijl} = T_{ij}^{-1} \sum_{t=1}^{T_{ij}} \{Y_{ijt} - \widehat{\mu}_j(U_{ijt})\} b_l(U_{ijt})$  and computed the sample variances of  $\widehat{\theta}_{ijl}$ , base on which performing the thresholding selection to encourage the sparsity. However, this approach is not applicable to sparsely observed functional data with bounded  $T_{ij}$ , as  $\theta_{ijl}$  can not be accurately estimated. We consider bridging the gap under the sparse case by applying the local linear smoothers to estimate the covariance functions  $\Sigma_{jj'}(u, v)$  directly. Note that  $\text{cov}(\theta_{ijl}, \theta_{ij'l'}) = \int b_l(u) \Sigma_{jj'}(u, v) b_{l'}(v) dudv$ . Instead of estimating  $\theta_{ijl}$ 's, we propose to directly estimate  $\text{cov}(\theta_{ijl}, \theta_{ij'l'})$  by  $\widehat{\text{cov}}(\theta_{ijl}, \theta_{ij'l'}) = \int_{\mathcal{U}} b_l(u) \widehat{\Sigma}_{jj'}(u, v) b_{l'}(v) dudv$ .

In the special case with  $j = j'$  and  $l = l'$ , this estimate degenerates to the estimated variance of  $\theta_{ijl}$ , and hence the thresholding idea proposed in Hu and Yao (2022) can still be employed. To establish the convergence properties under high-dimensional settings, it is essential to develop concentration results for  $\{\widehat{\text{cov}}(\theta_{jl}, \theta_{j'l'})\}$ . By the fact that

$$|\widehat{\text{cov}}(\theta_{ijl}, \theta_{ij'l'}) - \text{cov}(\theta_{ijl}, \theta_{ij'l'})| \leq \|\widehat{\Sigma}_{jj'} - \Sigma_{jj'}\|_{\mathcal{S}}, \quad (19)$$

our derived rate (11) in Theorem 7 becomes applicable. Specially, when  $j = j'$  and  $l = l'$ , (19), we can obtain the elementwise maximum rate for  $\widehat{\text{var}}(\theta_{ijl})$ . It is noteworthy that Hu and Yao (2022) relied on existing concentration inequalities for  $\chi_n^2$  to establish the concentration bound on the sample variance of  $\hat{\theta}_{ijt}$  under the dense design by assuming that  $\theta_{ijl}$  and  $\varepsilon_{ijt}$  are jointly Gaussian. By comparison, our proposal can handle both sparsely and densely observed functional data without requiring the Gaussianity assumption.

It is also worth mentioning that the above discussion also applies to other pre-fixed basis expansion methods when fitting functional additive regression models (Fan et al., 2014, 2015; Xue and Yao, 2021) or implementing functional linear discriminant analysis (Xue et al., 2024). Specifically, our proposed  $\{\widehat{\text{cov}}(\theta_{ijl}, \theta_{ij'l'})\}$  are also involved in the estimation, making Theorem 7 useful in this context.

### 4.3 Functional Thresholding

Our third application involves estimating the covariance matrix function  $\Sigma(\cdot, \cdot)$  in (1). Under the functional sparsity assumption that  $\Sigma$  belongs to a class of “approximately sparse” covariance matrix functions, Fang et al. (2024) proposed to perform adaptive functional thresholding on the local linear smoothers  $\{\widehat{\Sigma}_{jk}(\cdot, \cdot)\}_{j,k \in [p]}$  using entry-dependent functional thresholds that automatically adapt to the variability of  $\widehat{\Sigma}_{jk}(\cdot, \cdot)$ 's. Specifically, their adaptive functional thresholding estimator is defined as

$$\widetilde{\Sigma} = \{\widetilde{\Sigma}_{jk}(\cdot, \cdot)\}_{p \times p} \quad \text{with} \quad \widetilde{\Sigma}_{ij} = \Psi_{jk}^{1/2} \times s_{\lambda} \left( \frac{\widehat{\Sigma}_{jk}}{\Psi_{jk}} \right),$$

where, for any thresholding parameter  $\lambda \geq 0$ ,  $s_{\lambda}(\cdot)$  is the functional thresholding operator to enforce the functional sparsity with the aid of Hilbert–Schmidt norm, and  $\Psi_{jk}(\cdot, \cdot)$  is a surrogate estimator for the asymptotic variance of  $\widehat{\Sigma}_{jk}(\cdot, \cdot)$  (Fang et al., 2024). Alternatively, one can achieve a universal functional thresholding estimator

$$\check{\Sigma} = \{\check{\Sigma}_{jk}(\cdot, \cdot)\}_{p \times p} \quad \text{with} \quad \check{\Sigma}_{jk} = s_{\lambda}(\widehat{\Sigma}_{jk}),$$

where a universal threshold level is used for all entries. To investigate the theoretical properties of both functional thresholding estimators  $\widetilde{\Sigma}$  and  $\check{\Sigma}$  under high-dimension settings, it is crucial to make use of the elementwise maximum rate of  $\max_{j,k \in [p]} \|\widehat{\Sigma}_{jk} - \Sigma_{jk}\|_{\mathcal{S}}$ , as presented in Theorem 7. In contrast, Fang et al. (2024) assumed a rough rate (3) instead of providing a proof to facilitate their technical analysis.

Such functional sparsity assumption is restrictive for many data sets, especially in finance and economics, where the functional variables exhibit high correlations. To address this issue, Li et al. (2023) and Leng et al. (2024) employed the functional factor models framework for  $\mathbf{X}_i(\cdot)$ , which is decomposed as the sum of a common component driven by

low-dimensional latent factors and an idiosyncratic component  $\mathbf{e}_i(\cdot)$ . Instead of imposing the functional sparsity assumption on  $\Sigma$ , they imposed it on the covariance matrix function of  $\mathbf{e}_i(\cdot)$ , and proposed different estimators for  $\Sigma$  by performing the associated eigenanalysis to estimate the common covariance and then applying (adaptive) functional thresholding to the residual covariance. To develop the convergence rates of their proposed estimators, our elementwise maximum rate in Theorem 7 becomes applicable.

## 5. Simulations

In this section, we examine the finite-sample performance of the local linear smoothers for the mean and covariance function estimation in high dimensions.

We generalize the simulated example for univariate functional data in Zhang and Wang (2016) to the multivariate setting by generating

$$X_{ij}(u) = \mu_j(u) + \phi(u)^T \boldsymbol{\theta}_{ij}, \quad i \in [n], j \in [p], u \in \mathcal{U} = [0, 1],$$

where the true mean function  $\mu_j(u) = 1.5 \sin\{3\pi(u + 0.5)\} + 2u^3$ , the basis function  $\phi(u) = \{\sqrt{2} \cos(2\pi u), \sqrt{2} \sin(2\pi u), \sqrt{2} \cos(4\pi u), \sqrt{2} \sin(4\pi u)\}^T$  and the basis coefficient vector  $\boldsymbol{\theta}_i = (\boldsymbol{\theta}_{i1}^T, \dots, \boldsymbol{\theta}_{ip}^T)^T \in \mathbb{R}^{4p}$  is sampled independently from a mean zero multivariate Gaussian distribution with block covariance matrix  $\mathbf{\Lambda} \in \mathbb{R}^{4p \times 4p}$  whose  $(j, k)$ th block is given by  $\mathbf{\Lambda}_{jk} = \rho^{|j-k|} \text{diag}\{2^{-2}, \dots, 5^{-2}\} \in \mathbb{R}^{4 \times 4}$  for  $j, k \in [p]$ . Hence the  $(j, k)$ th entry of the true covariance functions  $\Sigma(\cdot, \cdot) = \{\Sigma_{jk}(\cdot, \cdot)\}_{p \times p}$  is  $\Sigma_{jk}(u, v) = \phi(u)^T \mathbf{\Lambda}_{jk} \phi(v)$ . We then generate the observed values  $Y_{ijt} = X_{ij}(U_{ijt}) + \varepsilon_{ijt}$  for  $t = 1, \dots, T_{ij} = T$ , where the time points  $U_{ijt}$ 's and errors  $\varepsilon_{ijt}$ 's are sampled independently from Uniform $[0, 1]$  and  $\mathcal{N}(0, 0.5^2)$ , respectively.

We use the Epanechnikov kernel with bandwidth values varying on a dense grid. To evaluate the performance of  $\hat{\mu}_j(\cdot)$  for  $j \in [p]$  and  $\hat{\Sigma}_{jk}(\cdot, \cdot)$  for  $(j, k) \in [p]^2$  given specific bandwidth  $h_{\mu,j}$  and  $h_{\Sigma,jk}$ , we define the corresponding mean integrated squared errors (MISE) as

$$\text{MISE}(\hat{\mu}_j, h_{\mu,j}) = \int \{\hat{\mu}_j(u) - \mu_j(u)\}^2 du, \quad \text{MISE}(\hat{\Sigma}_{jk}, h_{\Sigma,jk}) = \int \int \{\hat{\Sigma}_{jk}(u, v) - \Sigma_{jk}(u, v)\}^2 dudv.$$

We first calculate the elementwise minimal MISEs for the mean and covariance estimators over the grids of candidate bandwidths in prespecified sets  $\mathcal{H}_\mu$  and  $\mathcal{H}_\Sigma$ , respectively. We then compute their averages and maximums over  $j \in [p]$  for the mean functions, that is,

$$\begin{aligned} \text{AveMISE}(\mu) &= \frac{1}{p} \sum_j \min_{h_{\mu,j}} \text{MISE}(\hat{\mu}_j, h_{\mu,j}), \\ \text{MaxMISE}(\mu) &= \max_j \min_{h_{\mu,j}} \text{MISE}(\hat{\mu}_j, h_{\mu,j}), \end{aligned}$$

and over  $(j, k) \in [p]^2$  for the covariance functions, that is,

$$\begin{aligned} \text{AveMISE}(\Sigma) &= \frac{1}{p^2} \sum_j \sum_k \min_{h_{\Sigma,jk}} \text{MISE}(\hat{\Sigma}_{jk}, h_{\Sigma,jk}), \\ \text{MaxMISE}(\Sigma) &= \max_{j,k} \min_{h_{\Sigma,jk}} \text{MISE}(\hat{\Sigma}_{jk}, h_{\Sigma,jk}). \end{aligned}$$

We next use the example of estimating the mean functions to illustrate the rationale of the above measures. While  $\text{AveMISE}(\mu)$  presents the averaged elementwise minimal MISEs across  $j \in [p]$ , some simple calculations in Appendix B show that the attainable quantity of the minimal elementwise maximum of MISEs,  $\min_{(h_{\mu,1}, \dots, h_{\mu,p}) \in \mathcal{H}_\mu^p} \max_{j \in [p]} \text{MISE}(\hat{\mu}_j, h_{\mu,j})$ , is equal to  $\text{MaxMISE}(\mu)$ . It is worth noting that the optimal selection of bandwidths by minimizing, e.g., for mean functions, each  $\text{MISE}(\hat{\mu}_j, h_{\mu,j})$  over  $h_{\mu,j} \in \mathcal{H}_\mu$  or  $\max_j \text{MISE}(\hat{\mu}_j, h_{\mu,j})$  over  $(h_{\mu,1}, \dots, h_{\mu,p}) \in \mathcal{H}_\mu^p$  serves to validate our established theoretical results, assuming that  $\mu_j(\cdot)$ 's are known. In practical scenarios with unknown mean and covariance functions, one can employ the commonly-adopted cross-validation method to estimate the specific MISE as a function of bandwidths, whose minimizer produces the optimal bandwidth selection.




Figure 1: Plots of average MaxMISE (black) and AveMISE (red) against  $T$  with  $p = 50$  (solid), 100 (dashed) and 150 (dotted) for mean estimators (left) and covariance estimators (right).

We firstly consider settings of  $n = 100$ ,  $p = 50, 100, 150$ , and  $T = 5, 10, 20, 40, 60, 80, 100, 120, 140, 160, 180$ . We ran each simulation 100 times. Figure 1 plots the average AveMISE and MaxMISE as functions of  $T$  for the estimated mean and covariance functions. We observe that both MaxMISE and AveMISE display a similar trend as  $T$  increases with a steep decline followed by a slight decrease and then a period of stability. Such trend roughly corresponds to the three categories of “sparse”, “semi-dense”, and “ultra-dense”, respectively. In addition, while AveMISE reflects the performance for univariate functional data, MaxMISE gradually enlarges as  $p$  increases from 50 to 150, providing empirical evidence to support that the associated  $\log p$ -based convergence rates in high-dimensional settings.

As suggested by one referee, to further validate the established rates discussed in Remarks 5 and 8, we plot the average  $\log(\text{AveMISE})$  against  $\log n$  and the average  $\log(\text{MaxMISE})$  against  $\log(n/\log p)$  across different values of  $n$ ,  $p$  and  $T$  for estimated mean and covariance functions in Figure 2. Note that our theory indicates that  $\log(\text{AveMISE})$  and  $\log n$ , as well as  $\log(\text{MaxMISE})$  and  $\log(n/\log p)$ , both exhibit linear relationships, e.g., for sparse functional data, with a slope of  $-2/5$  for mean estimators and  $-1/3$  for covariance estimators. Furthermore, when the phase transitions from “semi-dense” to “ultra-dense” occur, they




Figure 2: Plots of average  $\log(\text{AveMISE})$  against  $\log n$  (left) and average  $\log(\text{MaxMISE})$  against  $\log(n/\log p)$  (right) for mean estimators (top) and covariance estimators (bottom) with  $p = 50, 100, 150$  and  $n = 50, 100, 150, 200, 250$ . The colored dashed lines correspond to different values of  $T$  ranging from 3 to 140, and the estimated slopes of the corresponding linear fits based on five points for  $\log n$  or fifteen points for  $\log(n/\log p)$  are also displayed. The slope of the black solid line presents the theoretical value  $-1/2$  (with the intercept being irrelevant here).

all admit linear relationships with a common slope of  $-1/2$ . Several apparent patterns are observable from Figure 2. First, both plots of  $\log(\text{AveMISE})$  against five values of  $\log n$  and  $\log(\text{MaxMISE})$  against fifteen values  $\log(n/\log p)$  show clear linear patterns across different values of  $T$ . This also demonstrates the  $\log p$ -based high-dimensional effect in elementwise maximum rates. Second, for sparse functional data (i.e.,  $T = 3, 5$ ), the estimated slopes of the linear fits are between  $-0.41$  and  $-0.37$  for mean estimators and between  $-0.37$  and  $-0.32$  for covariance estimators, which nicely align with the theoretical slope values of  $-2/5$  and  $-1/3$ , respectively. Third, as  $T$  grows from 10 to 140, the estimated slopes under all scenarios gradually increase to some values close to the theoretical slope value of  $-1/2$ , and

then stabilize, especially when measuring MaxMISE. This suggests that the phase transitions from “semi-dense” to “ultra-dense” may occur during the increase of  $T$ . For example, the slope for MaxMISE tends to be stable for  $T = 40, 60, 100, 140$ , indicating that the phase transition may occur around  $T = 40$  (earlier than the corresponding occurrence of phase transition for AveMISE). All of the above observations nicely validate our established theoretical results in Section 3.

### Acknowledgments

The authors would like to thank the Action Editor and three anonymous reviewers for their insightful comments and suggestions, which have led to significant improvement of our paper. Shaojun Guo was partially supported by the National Natural Science Foundation of China (No. 12471270). Dong Li was partially supported by the National Natural Science Foundation of China (No. 72471127). Xinghao Qiao was partially supported by the Seed Fund for Basic Research for New Staff at the University of Hong Kong.

## Appendix A. Technical Proofs

The appendix contains proofs of all theorems. Throughout, we use  $c, c_1, c_2, \dots$  to denote generic positive finite constants that may be different in different uses.

### A.1 Proof of Theorem 2

We organize the proof in four steps. First, we will define  $\hat{\mu}(\cdot), \tilde{\mu}(\cdot)$  and obtain the decomposition of  $\hat{\mu}(\cdot) - \tilde{\mu}(\cdot)$ . Second, we will prove the local concentration inequality for fixed interior point  $u \in \mathcal{U}$ . Third, we will prove the concentration inequality in  $L_2$  norm. Finally, we will prove the concentration inequality in supremum norm.

#### A.1.1 DEFINITION AND DECOMPOSITION

Without loss of generality, we let  $h_{\mu,j} = h$  for  $j \in [p]$  and denote  $\mathbf{e}_0 = (1, 0)^\top$ ,  $\tilde{\mathbf{U}}_{ijt} = \{1, (U_{ijt} - u)/h\}^\top$ ,

$$\begin{aligned}\hat{\mathbf{S}}_j(u) &= \sum_{i=1}^n v_{ij} \sum_{t=1}^{T_{ij}} \tilde{\mathbf{U}}_{ijt} \tilde{\mathbf{U}}_{ijt}^\top K_h(U_{ijt} - u), \\ \hat{\mathbf{R}}_j(u) &= \sum_{i=1}^n v_{ij} \sum_{t=1}^{T_{ij}} \tilde{\mathbf{U}}_{ijt} K_h(U_{ijt} - u) Y_{ijk}.\end{aligned}$$

A simple calculation yields that  $\hat{\mu}_j(u) = \mathbf{e}_0^\top \{\hat{\mathbf{S}}_j(u)\}^{-1} \hat{\mathbf{R}}_j(u)$ . Let

$$\tilde{\mu}_j(u) = \mathbf{e}_0^\top [\mathbb{E}\{\hat{\mathbf{S}}_j(u)\}]^{-1} \mathbb{E}\{\hat{\mathbf{R}}_j(u)\}. \quad (\text{A.1})$$

We can decompose  $\hat{\mu}_j(u) - \tilde{\mu}_j(u)$  as

$$\begin{aligned}\hat{\mu}_j(u) - \tilde{\mu}_j(u) &= \mathbf{e}_0^\top [\mathbb{E}\{\hat{\mathbf{S}}_j(u)\}]^{-1} [\hat{\mathbf{R}}_j(u) - \mathbb{E}\{\hat{\mathbf{R}}_j(u)\}] \\ &\quad - \mathbf{e}_0^\top \{\hat{\mathbf{S}}_j(u)\}^{-1} [\hat{\mathbf{S}}_j(u) - \mathbb{E}\{\hat{\mathbf{S}}_j(u)\}] [\mathbb{E}\{\hat{\mathbf{S}}_j(u)\}]^{-1} \hat{\mathbf{R}}_j(u),\end{aligned}$$

which then implies that

$$\begin{aligned} |\hat{\mu}_j(u) - \tilde{\mu}_j(u)| &\leq \|\mathbb{E}\{\hat{\mathbf{S}}_j(u)\}\|_{\min}^{-1} \|\hat{\mathbf{R}}_j(u) - \mathbb{E}\{\hat{\mathbf{R}}_j(u)\}\| \\ &\quad + \|\hat{\mathbf{S}}_j(u)\|_{\min}^{-1} \|\mathbb{E}\{\hat{\mathbf{S}}_j(u)\}\|_{\min}^{-1} \|\hat{\mathbf{R}}_j(u)\| \|\hat{\mathbf{S}}_j(u) - \mathbb{E}\{\hat{\mathbf{S}}_j(u)\}\|_{\text{F}}, \end{aligned} \quad (\text{A.2})$$

where, for any vector  $\mathbf{b} = (b_1, \dots, b_p)^\text{T}$ , we denote its Euclidean norm by  $\|\mathbf{b}\| = (\sum_i b_i^2)^{1/2}$  and, for any matrix  $\mathbf{B} = (B_{ij})_{p \times q}$ , we write  $\|\mathbf{B}\|_{\min} = \{\lambda_{\min}(\mathbf{B}^\text{T}\mathbf{B})\}^{1/2}$  and  $\|\mathbf{B}\|_{\text{F}} = (\sum_{i,j} B_{ij}^2)^{1/2}$  to denote its Frobenius norm.

### A.1.2 LOCAL CONCENTRATION INEQUALITY

We will firstly show that there exists some positive constant  $c$  (independent of  $u$ ) such that for any  $\delta > 0$  and  $u \in \mathcal{U}$ ,

$$\mathbb{P}\left\{\|\hat{\mathbf{S}}_j(u) - \mathbb{E}\{\hat{\mathbf{S}}_j(u)\}\|_{\text{F}} \geq \delta\right\} \leq 8 \exp\left(-\frac{cn\bar{T}_{\mu,j}h\delta^2}{1+\delta}\right). \quad (\text{A.3})$$

For  $k, l = 1, 2$ , let  $\hat{S}_{jkl}(u)$  be the  $(k, l)$ th entry of  $\hat{\mathbf{S}}_j(u)$ . Under Assumptions 4 and 6, we obtain that for any integer  $q = 2, 3, \dots$  and  $s = 0, 1, 2$ ,

$$\mathbb{E}\left\{\left|\left(\frac{U_{ijt} - u}{h}\right)^s K_h(U_{ijt} - u)\right|^q\right\} \leq \int h^{-q} K^q\left(\frac{t-u}{h}\right) \left|\frac{t-u}{h}\right|^{sq} f_U(t) dt \leq ch^{1-q}. \quad (\text{A.4})$$

Note that Assumption 3 implies that the weights  $v_{ij}$ 's are of the same order  $v_{ij} \asymp (n\bar{T}_{\mu,j})^{-1}$ . By (A.4), it holds that

$$\begin{aligned} \sum_{i=1}^n \sum_{t=1}^{T_{ij}} \mathbb{E}\left\{\left|\left(\frac{U_{ijt} - u}{h}\right)^s K_h(U_{ijt} - u)\right|^2\right\} &\leq cn\bar{T}_{\mu,j}h^{-1}, \\ \sum_{i=1}^n \sum_{t=1}^{T_{ij}} \mathbb{E}\left\{\left|\left(\frac{U_{ijt} - u}{h}\right)^s K_h(U_{ijt} - u)\right|^q\right\} &\leq 2^{-1}q!cn\bar{T}_{\mu,j}h^{-1}h^{2-q} \quad \text{for } q \geq 3. \end{aligned}$$

By the Bernstein inequality (see Theorem 2.10 and Corollary 2.11 of Boucheron et al., 2013), we obtain that there exists some positive constant  $c$  (independent of  $u$ ) such that for any  $\delta > 0$  and  $u \in \mathcal{U}$ ,

$$\mathbb{P}\left\{|\hat{S}_{jkl}(u) - \mathbb{E}\{\hat{S}_{jkl}(u)\}| \geq \delta\right\} \leq 2 \exp\left(-\frac{cn\bar{T}_{\mu,j}h\delta^2}{1+\delta}\right),$$

for  $k, l = 1, 2$ , which, by the union bound of probability, implies that (A.3) holds.

For  $k = 1, 2$ , let  $\hat{R}_{jk}(u)$  be the  $k$ th element of  $\hat{\mathbf{R}}_j(u)$ . We will next show that there exists some positive constant  $c$  (independent of  $u$ ) such that for any  $\delta > 0$  and  $u \in \mathcal{U}$ ,

$$\mathbb{P}\left\{|\hat{R}_{jk}(u) - \mathbb{E}\{\hat{R}_{jk}(u)\}| \geq \delta\right\} \leq c \exp\left(-\frac{c\gamma_{n,T,h,j}\delta^2}{1+\delta}\right), \quad (\text{A.5})$$

where  $\gamma_{n,T,h,j} = n(1 \wedge \bar{T}_{\mu,j}h)$ . We only need to consider the case  $k = 1$ , while the case  $k = 2$  can be demonstrated in a similar manner. Denote that

$$\begin{aligned}\xi_{ijt} &= Y_{ijt} - \mu_j(U_{ijt}), \\ \hat{R}_{j3}(u) &= \sum_{i=1}^n v_{ij} \sum_{t=1}^{T_{ij}} K_h(U_{ijt} - u) \mu_j(U_{ijt}), \\ \hat{R}_{j4}(u) &= \sum_{i=1}^n v_{ij} \sum_{t=1}^{T_{ij}} K_h(U_{ijt} - u) \xi_{ijt}.\end{aligned}$$

Then  $\hat{R}_{j1}(u) - \mathbb{E}\{\hat{R}_{j1}(u)\}$  can be rewritten as

$$\hat{R}_{j1}(u) - \mathbb{E}\{\hat{R}_{j1}(u)\} = \hat{R}_{j3}(u) - \mathbb{E}\{\hat{R}_{j3}(u)\} + \hat{R}_{j4}(u). \quad (\text{A.6})$$

Following the same procedure to prove (A.3) and using the Bernstein inequality, we can obtain that there exists some positive constant  $c$  such that for any  $\delta > 0$  and  $u \in \mathcal{U}$ ,

$$\mathbb{P}\left\{|\hat{R}_{j3}(u) - \mathbb{E}\{\hat{R}_{j3}(u)\}| \geq \delta\right\} \leq 2 \exp\left(-\frac{cn\bar{T}_{\mu,j}h\delta^2}{1+\delta}\right). \quad (\text{A.7})$$

Now we consider the tail behavior of  $\hat{R}_{j4}(u)$ . Define the event  $V_j = \{U_{ijt}, t \in [T_{ij}], i \in [n]\}$ . Rewrite  $\hat{R}_{j4}(u) = \sum_{i=1}^n v_{ij} \psi_{ij1}(u)$  with  $\psi_{ij1}(u) = \frac{\sum_{t=1}^{T_{ij}} K_h(U_{ijt}-u) \xi_{ijt}}{\sum_{t=1}^{T_{ij}} K_h(U_{ijt}-u)}$ . If  $\sum_{t=1}^{T_{ij}} K_h(U_{ijt}-u) > 0$ , by Jensen's inequality, we have

$$\begin{aligned}& \mathbb{E}\left[\exp\{\lambda v_{ij} \psi_{ij1}(u)\} \middle| V_j\right] \\ &= \mathbb{E}\left(\exp\left[\lambda v_{ij} \frac{\{\sum_{t=1}^{T_{ij}} K_h(U_{ijt}-u) \xi_{ijt}\} \{\sum_{t=1}^{T_{ij}} K_h(U_{ijt}-u)\}}{\sum_{t=1}^{T_{ij}} K_h(U_{ijt}-u)}\right] \middle| V_j\right) \\ &\leq \frac{1}{\sum_{t=1}^{T_{ij}} K_h(U_{ijt}-u)} \sum_{t=1}^{T_{ij}} K_h(U_{ijt}-u) \mathbb{E}\left[\exp\left\{\lambda v_{ij} \xi_{ijt} \sum_{t'=1}^{T_{ij}} K_h(U_{ijt'-u})\right\} \middle| V_j\right] \\ &\leq \exp\left[\lambda^2 c^2 (n\bar{T}_{\mu,j})^{-2} \left\{\sum_{t=1}^{T_{ij}} K_h(U_{ijt}-u)\right\}^2\right].\end{aligned}$$

Note that the last line comes from the fact that  $\mathbb{E}\{\exp(\lambda \xi_{ijt}) | V_j\} \leq \exp(\lambda^2 c^2)$  for any  $\lambda \in \mathbb{R}$ , which is implied from the sub-Gaussianities in Assumption 1 and (4). Clearly, the above inequality still holds even if  $\sum_{t=1}^{T_{ij}} K_h(U_{ijt}-u) = 0$ . Assumption 5 implies that the number of nonzero terms in  $\sum_{t=1}^{T_{ij}} K_h(U_{ijt}-u)$  has an upper bound  $c(1 \vee \bar{T}_{\mu,j}h)$ , which yields that

$\sum_{t=1}^{T_{ij}} K_h(U_{ijt} - u) \leq ch^{-1}(1 \vee \bar{T}_{\mu,j}h)$ . Therefore, for any  $\lambda \in \mathbb{R}$ , we obtain that

$$\begin{aligned} \mathbb{E} \left[ \exp \left\{ \lambda \sum_{i=1}^n v_{ij} \psi_{ij1}(u) \right\} \middle| V_j \right] &= \prod_{i=1}^n \mathbb{E} \left[ \exp \left\{ \lambda v_{ij} \psi_{ij1}(u) \right\} \middle| V_j \right] \\ &\leq \exp \left[ \lambda^2 c^2 \sum_{i=1}^n v_{ij}^2 \left\{ \sum_{t=1}^{T_{ij}} K_h(U_{ijt} - u) \right\}^2 \right] \\ &\leq \exp \left\{ \lambda^2 c^2 (n \bar{T}_{\mu,j} h)^{-2} h (1 \vee \bar{T}_{\mu,j} h) \sum_{i=1}^n \sum_{t=1}^{T_{ij}} K_h(U_{ijt} - u) \right\}. \end{aligned}$$

For any  $\delta > 0$ , define the event  $\Omega_{j,1}(\delta) = \left\{ \sum_{i=1}^n \sum_{t=1}^{T_{ij}} K_h(U_{ijt} - u) \leq c(1 + \delta)n\bar{T}_{\mu,j} \right\}$ . We have

$$\begin{aligned} \mathbb{E} \left[ \exp \left\{ \lambda \sum_{i=1}^n v_{ij} \psi_{ij1}(u) \right\} \middle| \Omega_{j,1}(\delta) \right] &\leq \exp \left\{ \lambda^2 c^2 (1 + \delta) (n \bar{T}_{\mu,j} h)^{-2} n \bar{T}_{\mu,j} h (1 \vee \bar{T}_{\mu,j} h) \right\} \\ &= \exp \left\{ (1 + \delta) \lambda^2 c^2 (n \bar{T}_{\mu,j} h)^{-1} (1 \vee \bar{T}_{\mu,j} h) \right\}. \end{aligned}$$

As a consequence, we obtain that

$$\mathbb{P} \left\{ \sum_{i=1}^n v_{ij} \psi_{ij1}(u) \geq \delta \middle| \Omega_{j,1}(\delta) \right\} \leq \exp \left\{ -\lambda \delta + (1 + \delta) \lambda^2 c^2 (n \bar{T}_{\mu,j} h)^{-1} (1 \vee \bar{T}_{\mu,j} h) \right\}. \quad (\text{A.8})$$

With the choice of  $\lambda = n \bar{T}_{\mu,j} h \delta / \{2(1 + \delta)c^2(1 \vee \bar{T}_{\mu,j}h)\}$ , (A.8) degenerates to

$$\mathbb{P} \left\{ \sum_{i=1}^n v_{ij} \psi_{ij1}(u) \geq \delta \middle| \Omega_{j,1}(\delta) \right\} \leq \exp \left\{ -\frac{cn(1 \wedge \bar{T}_{\mu,j}h)\delta^2}{1 + \delta} \right\}. \quad (\text{A.9})$$

Note that  $\sum_{i=1}^n \sum_{t=1}^{T_{ij}} \mathbb{E}\{K_h(U_{ijt} - u)\} \leq cn\bar{T}_{\mu,j}$ . By the Bernstein inequality, we obtain that there exists some positive constant  $c$  such that for any  $\delta > 0$ ,

$$\mathbb{P} \left( \sum_{i=1}^n \sum_{t=1}^{T_{ij}} \left[ K_h(U_{ijt} - u) - \mathbb{E}\{K_h(U_{ijt} - u)\} \right] \geq n \bar{T}_{\mu,j} \delta \right) \leq \exp \left( -\frac{cn \bar{T}_{\mu,j} h \delta^2}{1 + \delta} \right),$$

which implies that

$$1 - \mathbb{P}\{\Omega_{j,1}(\delta)\} \leq \exp \left( -\frac{cn \bar{T}_{\mu,j} h \delta^2}{1 + \delta} \right). \quad (\text{A.10})$$

Combining (A.9) and (A.10), we obtain that there exists some constant  $c > 0$  such that for any  $\delta > 0$ ,

$$\mathbb{P} \left\{ \widehat{R}_{j4}(u) \geq \delta \right\} \leq \mathbb{P} \left\{ \widehat{R}_{j4}(u) \geq \delta \middle| \Omega_{j,1}(\delta) \right\} + \mathbb{P} \left\{ \Omega_{j,1}(\delta)^c \right\} \leq 2 \exp \left\{ -\frac{cn(1 \wedge \bar{T}_{\mu,j}h)\delta^2}{1 + \delta} \right\},$$

and, consequently,

$$\mathbb{P}\left\{|\widehat{R}_{j4}(u)| \geq \delta\right\} \leq 4 \exp\left\{-\frac{cn(1 \wedge \bar{T}_{\mu,j}h)\delta^2}{1+\delta}\right\}.$$

This together with (A.6) and (A.7) yields that, there exists some constant  $c$  such that

$$\mathbb{P}\left\{|\widehat{R}_{j1}(u) - \mathbb{E}\{\widehat{R}_{j1}(u)\}| \geq \delta\right\} \leq 6 \exp\left\{-\frac{cn(1 \wedge \bar{T}_{\mu,j}h)\delta^2}{1+\delta}\right\}.$$

Define the event  $\Omega_{j,2}(\delta) = \{\|\widehat{\mathbf{S}}_j(u) - \mathbb{E}\{\widehat{\mathbf{S}}_j(u)\}\|_{\mathbb{F}} \leq \delta/2\}$ . Note that  $\mathbb{E}\{\widehat{\mathbf{S}}_j(u)\}$  is positive definite. On the event  $\Omega_{j,2}(\delta)$  with  $\delta \in (0, 1]$ , we obtain that

$$\|\widehat{\mathbf{S}}_j(u)\|_{\min} \geq c(1 - \delta/2). \quad (\text{A.11})$$

By (A.3), we have

$$1 - \mathbb{P}\{\Omega_{j,2}(\delta)\} \leq 8 \exp\left(-\frac{cn\bar{T}_{\mu,j}h\delta^2}{1+\delta}\right). \quad (\text{A.12})$$

Define the event  $\Omega_{j,3}(\delta) = \{\|\widehat{\mathbf{R}}_j(u) - \mathbb{E}\{\widehat{\mathbf{R}}_j(u)\}\| \leq \delta\}$ . Note that, under Assumption 3 with  $v_{ij} = (n\bar{T}_{\mu,j})^{-1}$ ,  $\sum_{i=1}^n v_{ij} \sum_{t=1}^{T_{ij}} \mathbb{E}\{K_h(U_{ijt} - u)\} \leq c$  and  $\mu_j(\cdot)$  is uniformly bounded over  $\mathcal{U}$ , hence  $\|\mathbb{E}\{\widehat{\mathbf{R}}_j(u)\}\|$  is uniformly bounded over  $\mathcal{U}$ . On the event  $\Omega_{j,3}(\delta)$ , we have

$$\|\widehat{\mathbf{R}}_j(u)\| \leq c(1 + \delta). \quad (\text{A.13})$$

On the event  $\Omega_{j,2}(\delta) \cap \Omega_{j,3}(\delta)$  with  $\delta \in (0, 1]$ , it follows from (A.2), (A.11) and (A.13) that

$$|\widehat{\mu}_j(u) - \widetilde{\mu}_j(u)| \leq c\delta + c(1 - \delta/2)^{-1}(1 + \delta)\delta \leq c_3\delta.$$

This together with concentration inequalities in (A.5) and (A.12) implies that there exist some positive universal constants  $c_1$  and  $c_2$  such that for any  $\delta \in (0, 1]$  and  $u \in \mathcal{U}$ ,

$$\mathbb{P}\left\{|\widehat{\mu}_j(u) - \widetilde{\mu}_j(u)| \geq \delta\right\} \leq c_2 \exp(-c_1\gamma_{n,T,h,j}\delta^2),$$

which completes the proof of local concentration inequality for the mean estimator.

### A.1.3 CONCENTRATION INEQUALITY IN $L_2$ NORM

In the proof, we need the following lemma.

**Lemma 11** *Let  $X$  be a random variable. If for some constants  $c_1, c_2 > 0$ ,  $\mathbb{P}(|X| > \delta) \leq c_1 \exp\{-c_2^{-1} \min(\delta^2, \delta)\}$  for any  $\delta > 0$ , then for any integer  $q \geq 1$ ,*

$$\mathbb{E}(X^{2q}) \leq q!c_1(4c_2)^q + (2q)!c_1(4c_2)^{2q}.$$

*Conversely, if for some positive constants  $a_1, a_2$ ,  $\mathbb{E}(X^{2q}) \leq q!a_1a_2^q + (2q)!a_1a_2^{2q}$  for any integer  $q \geq 1$ , then by letting  $c_1^* = a_1$  and  $c_2^* = 32(a_2 + a_2^2)$ , we have that*

$$\mathbb{P}(|X| > \delta) \leq c_1^* \exp\{-c_2^{*-1} \min(\delta^2, \delta)\}$$

for any  $\delta > 0$ .

**Proof** This lemma can be proved in a similar way to Theorem 2.3 of Boucheron et al. (2013) and hence the proof is omitted here. In the proof, the following two inequalities are used, i.e., for any  $c, \delta > 0$ ,

$$\frac{1}{2} \min(\delta^2, \delta) \leq \frac{\delta^2}{1 + \delta} \leq \min(\delta^2, \delta)$$

and

$$\sqrt{\frac{c\delta}{2}} + \frac{c\delta}{2} \leq \frac{c(\delta + \sqrt{\delta^2 + 4\delta/c})}{2} \leq \sqrt{c\delta} + c\delta. \quad \blacksquare$$

We are now ready to derive the  $L_2$  concentration inequality of  $\|\hat{\mu}_j - \tilde{\mu}_j\|_2$ . Let

$$\tilde{\mathbf{S}}_j(u) = (n\bar{T}_{\mu,j})^{-1} \sum_{i=1}^n \sum_{t=1}^{T_{ij}} \tilde{\mathbf{U}}_{ijt} \tilde{\mathbf{U}}_{ijt}^T K_h(U_{ijt} - u). \quad (\text{A.14})$$

Then we have that  $\|\hat{\mathbf{S}}_j(u)\|_{\min} \geq c\|\tilde{\mathbf{S}}_j(u)\|_{\min}$ . We now give a lower bound on  $\|\tilde{\mathbf{S}}_j(u)\|_{\min}$ . Denote  $W = \sup_{u \in \mathcal{U}} \|\tilde{\mathbf{S}}_j(u) - \mathbb{E}\{\tilde{\mathbf{S}}_j(u)\}\|_{\text{F}}$ . Let  $\tilde{S}_{jkl}(u)$  be the  $(k, l)$ th entry of  $\tilde{\mathbf{S}}_j(u)$  for  $k, l = 1, 2$ . Note that  $\mathbb{E}\{|(U_{ijt} - u)^a h^{-a} K_h(U_{ijt} - u)|\} \leq c$  and  $\mathbb{E}(W) \leq 4 \max_{k,l} \mathbb{E}\{\sup_{u \in \mathcal{U}} |\tilde{S}_{jkl}(u)|\}$  for  $a = 0, 1, 2$ . In an analogy to Lemma 13.5 of Boucheron et al. (2013), we can show that  $\mathbb{E}(W) \leq c(n\bar{T}_{\mu,j})^{-1/2}$ . Note that Lemma 13.5 of Boucheron et al. (2013) relies on the results presented in Lemma 13.1 of Boucheron et al. (2013). Consequently, Lemma 13.5 assumes that the corresponding index set is countable in order to apply Lemma 13.1, as the supremums of the summation of indicator functions may not be measurable. However, in our specific case, each component of  $\tilde{\mathbf{S}}_j(u) - \mathbb{E}\{\tilde{\mathbf{S}}_j(u)\}$  is the sum of continuous functions, for which the supremums over  $\mathcal{U}$  are measurable. Therefore, when we extend the index set from countable to the uncountable set  $\mathcal{U}$ , this lemma, as well as Theorems 11.10 and 12.5 of Boucheron et al. (2013), still hold true and can be applicable to our situation. Moreover, it follows from the facts  $\text{var}(W) \leq \mathbb{E}(W^2) \leq 4 \max_{k,l} \text{var}\{\sup_{u \in \mathcal{U}} \tilde{S}_{jkl}(u)\}$ ,  $|(U_{ijt} - u)^a h^{-a} K_h(U_{ijt} - u)| \leq ch^{-1}$ ,  $\mathbb{E}\{(U_{ijt} - u)^{2a} h^{-2a} K_h^2(U_{ijt} - u)\} \leq ch^{-1}$  for  $a = 0, 1, 2$ , and Theorem 11.10 of Boucheron et al. (2013) that  $\text{var}(n\bar{T}_{\mu,j}hW) \leq 2\mathbb{E}(n\bar{T}_{\mu,j}hW) + \sum_{i=1}^n \sum_{t=1}^{T_{ij}} ch^{-1}h^2 \leq c(n\bar{T}_{\mu,j})^{1/2}h + cn\bar{T}_{\mu,j}h$ , which implies that the variance of  $W$  is bounded by  $c(n\bar{T}_{\mu,j}h)^{-1} \leq c\gamma_{n,T,h,j}^{-1}$ . Applying Theorem 12.5 of Boucheron et al. (2013) yields that there exists some positive constant  $c$  such that, for any  $\delta > 0$ ,

$$\mathbb{P}\{W - \mathbb{E}(W) > \delta\} \leq \exp\left(-\frac{c\gamma_{n,T,h,j}\delta^2}{1 + \delta}\right). \quad (\text{A.15})$$

Define the event  $\Omega_{j,4}(\delta) = \{\sup_{u \in \mathcal{U}} \|\tilde{\mathbf{S}}_j(u) - \mathbb{E}\{\tilde{\mathbf{S}}_j(u)\}\|_{\text{F}} \leq \delta/2\}$  with  $\delta \in (0, 1]$ . By (A.15), we obtain that there exists some constant  $c > 0$  such that, for any  $\delta \in (0, 1]$ ,

$$1 - \mathbb{P}\{\Omega_{j,4}(\delta)\} \leq 2 \exp(-c\gamma_{n,T,h,j}\delta^2). \quad (\text{A.16})$$

On the event  $\Omega_{j,4} = \Omega_{j,4}(\delta_1)$  with  $c\gamma_{n,T,h,j}^{-1/2} < \delta_1 \leq 1$ ,  $\|\hat{\mathbf{S}}_j(u)\|_{\min} \geq c\|\tilde{\mathbf{S}}_j(u)\|_{\min} \geq c(1 - \delta_1/2) \geq c/2$ . Note that  $\mathbb{E}\{\hat{\mathbf{S}}_j(u)\}$  is positive definite and  $\|\mathbb{E}\{\hat{\mathbf{R}}_j(u)\}\|$  is uniformly bounded

over  $\mathcal{U}$ . On the event  $\Omega_{j,4}$ , it thus follows from (A.2) and  $\|\widehat{\mathbf{R}}_j(u)\| \leq \|\widehat{\mathbf{R}}_j(u) - \mathbb{E}\{\widehat{\mathbf{R}}_j(u)\}\| + \|\mathbb{E}\{\widehat{\mathbf{R}}_j(u)\}\|$  that

$$|\widehat{\mu}_j(u) - \widetilde{\mu}_j(u)| \leq c\|\widehat{\mathbf{R}}_j(u) - \mathbb{E}\{\widehat{\mathbf{R}}_j(u)\}\| + c\|\widehat{\mathbf{S}}_j(u) - \mathbb{E}\{\widehat{\mathbf{S}}_j(u)\}\|_{\mathbb{F}}, \quad (\text{A.17})$$

where the positive constant  $c$  does not depend on  $u \in \mathcal{U}$ .

Combining (A.17) with (A.3), (A.5) and applying the first part of Lemma 11 yields that, for any  $u \in \mathcal{U}$  and integer  $q \geq 1$ ,

$$\mathbb{E}\left\{|\widehat{\mu}_j(u) - \widetilde{\mu}_j(u)|^{2q} \middle| \Omega_{j,4}\right\} \leq q!c\left(\frac{4}{c\gamma_{n,T,h,j}}\right)^q + (2q)!c\left(\frac{4}{c\gamma_{n,T,h,j}}\right)^{2q}.$$

Applying the second part of Lemma 11 and (A.16), we can show that, for each  $\delta \in (0, 1]$ ,

$$\mathbb{P}(\|\widehat{\mu}_j - \widetilde{\mu}_j\|_2 \geq \delta) \leq \mathbb{P}(\|\widehat{\mu}_j - \widetilde{\mu}_j\|_2 \geq \delta | \Omega_{j,4}) + \mathbb{P}(\Omega_{j,4}^c) \leq c_2 \exp(-c_1\gamma_{n,T,h,j}\delta^2),$$

which means that (5) in Theorem 2 holds and completes the proof of concentration inequality for the mean estimator in  $L_2$  norm.

#### A.1.4 CONCENTRATION INEQUALITY IN SUPREMUM NORM

We will derive the uniform concentration bound of  $\sup_{u \in \mathcal{U}} |\widehat{\mu}_j(u) - \widetilde{\mu}_j(u)|$ . We partition the interval  $\mathcal{U} = [0, 1]$  into  $N$  subintervals  $I_s$  for  $s \in [N]$  of equal length. Let  $u_s$  be the center of  $I_s$ , then we have

$$\sup_{u \in \mathcal{U}} |\widehat{\mu}_j(u) - \widetilde{\mu}_j(u)| \leq \max_{s \in [N]} \left[ |\widehat{\mu}_j(u_s) - \widetilde{\mu}_j(u_s)| + \left| \{\widehat{\mu}_j(u_s) - \widehat{\mu}_j(u)\} - \{\widetilde{\mu}_j(u_s) - \widetilde{\mu}_j(u)\} \right| \right].$$

We need to bound the second term. By some calculations, it suffices to bound  $\left| \{\widehat{R}_{jk}(u) - \widehat{R}_{jk}(u_s)\} - [\mathbb{E}\{\widehat{R}_{jk}(u)\} - \mathbb{E}\{\widehat{R}_{jk}(u_s)\}] \right|$  and  $\left| \{\widehat{S}_{jkl}(u) - \widehat{S}_{jkl}(u_s)\} - [\mathbb{E}\{\widehat{S}_{jkl}(u)\} - \mathbb{E}\{\widehat{S}_{jkl}(u_s)\}] \right|$  for  $k, l = 1, 2$ , which means that we need to bound  $|\widehat{R}_{jk}(u) - \widehat{R}_{jk}(u_s)|$  and  $|\widehat{S}_{jkl}(u) - \widehat{S}_{jkl}(u_s)|$ . Let  $u \in I_s$  and consider  $|\widehat{R}_{j1}(u) - \widehat{R}_{j1}(u_s)|$  first. Define the event  $\Omega_{R,j1} = \left\{ \sum_{i=1}^n v_{ij} \sum_{t=1}^{T_{ij}} |Y_{ijt}| \leq \mathbb{E}(\sum_{i=1}^n v_{ij} \sum_{t=1}^{T_{ij}} |Y_{ijt}|) + 1 \right\}$ . On this event, it follows from Assumption 6(ii) that

$$\begin{aligned} |\widehat{R}_{j1}(u) - \widehat{R}_{j1}(u_s)| &\leq \left| \sum_{i=1}^n v_{ij} \sum_{t=1}^{T_{ij}} Y_{ijt} \{K_h(U_{ijt} - u) - K_h(U_{ijt} - u_s)\} \right| \\ &\leq \frac{c|u - u_s|}{h^2} \sum_{i=1}^n v_{ij} \sum_{t=1}^{T_{ij}} |Y_{ijt}| \leq \frac{c}{Nh^2} \left\{ \mathbb{E} \left( \sum_{i=1}^n v_{ij} \sum_{t=1}^{T_{ij}} |Y_{ijt}| \right) + 1 \right\} \leq \frac{c}{Nh^2}. \end{aligned}$$

Applying similar techniques as above, we can define events  $\Omega_{R,jk}$  and  $\Omega_{S,jkl}$  for  $k, l = 1, 2$ . On the intersection of these events, we can obtain that  $|\widehat{R}_{jk}(u) - \widehat{R}_{jk}(u_s)| \leq c(Nh^2)^{-1}$  and  $|\widehat{S}_{jkl}(u) - \widehat{S}_{jkl}(u_s)| \leq c(Nh^2)^{-1}$ . Combing the above results, we have

$$\sup_{u \in \mathcal{U}} |\widehat{\mu}_j(u) - \widetilde{\mu}_j(u)| \leq \max_{s \in [N]} |\widehat{\mu}_j(u_s) - \widetilde{\mu}_j(u_s)| + \frac{c}{Nh^2}.$$

Applying Hoeffding's inequality, we obtain that  $\mathbb{P}(\Omega_{R,jk}^c) \leq \exp\{-(2 \sum_{i=1}^n c^2 v_{ij}^2 T_{ij}^2)^{-1}\} = \exp(-cn) \leq \exp(-c\gamma_{n,T,h,j})$  and  $\mathbb{P}(\Omega_{S,jkl}^c) \leq \exp\{-(2 \sum_{i=1}^n c^2 v_{ij}^2 T_{ij}^2)^{-1}\} = \exp(-cn) \leq \exp(-c\gamma_{n,T,h,j})$  for  $k, l = 1, 2$ . It follows from the above results and the union bound of probability with the choice of  $N = \lfloor c(h^2\delta)^{-1} \rfloor$  that there exist some positive constants  $c_1$  and  $c_2$  such that, for any  $\delta \in (0, 1]$ ,

$$\mathbb{P}\left\{\sup_{u \in \mathcal{U}} |\hat{\mu}_j(u) - \tilde{\mu}_j(u)| \geq \delta\right\} \leq \frac{c_2}{h^2\delta} \exp(-c_1\gamma_{n,T,h,j}\delta^2). \quad (\text{A.18})$$

Take arbitrarily small  $\epsilon_1 > 0$ . If  $n^{\epsilon_1}\gamma_{n,T,h,j}\delta^2 \geq 1$ , then the right side of (A.18) reduces to  $c_2\{n^{\epsilon_1}\gamma_{n,T,h,j}\}^{1/2}h^{-2} \exp(-c_1\gamma_{n,T,h,j}\delta^2)$ . If  $n^{\epsilon_1}\gamma_{n,T,h,j}\delta^2 \leq 1$ , we can choose  $c_2$  and  $n^{\epsilon_1} > c$  such that  $c_2 \exp(-c_1c^{-1}) \geq 1$  and the same bound  $c_1\{n^{\epsilon_1}\gamma_{n,T,h,j}\}^{1/2}h^{-2} \exp(-c_1\gamma_{n,T,h,j}\delta^2)$  can still be used. Hence (6) in Theorem 2 holds, which completes the proof of concentration inequality for the mean estimator in supremum norm.  $\blacksquare$

## A.2 Proof of Theorem 3

We organize the proof in four steps. First, we will define  $\hat{\Sigma}(\cdot, \cdot)$ ,  $\tilde{\Sigma}(\cdot, \cdot)$  and obtain the decomposition of  $\hat{\Sigma}(\cdot, \cdot) - \tilde{\Sigma}(\cdot, \cdot)$ . Second, we will prove the local concentration inequality for fixed  $(u, v) \in \mathcal{U}^2$ . Third, we will prove the concentration inequality in Hilbert-Schmidt norm. Finally, we will prove the concentration inequality in supremum norm.

### A.2.1 DEFINITION AND DECOMPOSITION

Without loss of generality, let  $h_{\Sigma,jk} = h$  for  $(j, k) \in [p]^2$  and denote  $\tilde{\mathbf{e}}_0 = (1, 0, 0)^\top$ ,  $\tilde{\mathbf{U}}_{ijjts} = \{1, (U_{ijt} - u)/h, (U_{iks} - v)/h\}^\top$ . For  $j = k$ , let

$$\begin{aligned} \hat{\mathbf{\Xi}}_{jj}(u, v) &= \sum_{i=1}^n w_{ijj} \sum_{1 \leq t \neq s \leq T_{ij}} \tilde{\mathbf{U}}_{ijjts} \tilde{\mathbf{U}}_{ijjts}^\top K_h(U_{ijt} - u) K_h(U_{ijs} - v), \\ \hat{\mathbf{Z}}_{jj}(u, v) &= \sum_{i=1}^n w_{ijj} \sum_{1 \leq t \neq s \leq T_{ij}} \tilde{\mathbf{U}}_{ijjts} \Theta_{ijjts} K_h(U_{ijt} - u) K_h(U_{ijs} - v). \end{aligned}$$

For  $j \neq k$ , let

$$\begin{aligned} \hat{\mathbf{\Xi}}_{jk}(u, v) &= \sum_{i=1}^n w_{ijk} \sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} \tilde{\mathbf{U}}_{ijjts} \tilde{\mathbf{U}}_{ijjts}^\top K_h(U_{ijt} - u) K_h(U_{iks} - v), \\ \hat{\mathbf{Z}}_{jk}(u, v) &= \sum_{i=1}^n w_{ijk} \sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} \tilde{\mathbf{U}}_{ijjts} \Theta_{ijjts} K_h(U_{ijt} - u) K_h(U_{iks} - v). \end{aligned}$$

A simple calculation yields that  $\hat{\Sigma}_{jk}(u, v) = \tilde{\mathbf{e}}_0^\top \{\hat{\mathbf{\Xi}}_{jk}(u, v)\}^{-1} \hat{\mathbf{Z}}_{jk}(u, v)$ . Let

$$\tilde{\Sigma}_{jk}(u, v) = \tilde{\mathbf{e}}_0^\top [\mathbb{E}\{\hat{\mathbf{\Xi}}_{jk}(u, v)\}]^{-1} \mathbb{E}\{\hat{\mathbf{Z}}_{jk}(u, v)\}. \quad (\text{A.19})$$

We can decompose  $\hat{\Sigma}_{jk}(u, v) - \tilde{\Sigma}_{jk}(u, v)$  as

$$\begin{aligned} \hat{\Sigma}_{jk}(u, v) - \tilde{\Sigma}_{jk}(u, v) &= \tilde{\mathbf{e}}_0^\top (\{\hat{\mathbf{\Xi}}_{jk}(u, v)\}^{-1} - [\mathbb{E}\{\hat{\mathbf{\Xi}}_{jk}(u, v)\}]^{-1}) \hat{\mathbf{Z}}_{jk}(u, v) \\ &\quad + \tilde{\mathbf{e}}_0^\top [\mathbb{E}\{\hat{\mathbf{\Xi}}_{jk}(u, v)\}]^{-1} [\hat{\mathbf{Z}}_{jk}(u, v) - \mathbb{E}\{\hat{\mathbf{Z}}_{jk}(u, v)\}], \end{aligned}$$

which further implies that

$$\begin{aligned}
 & \left| \widehat{\Sigma}_{jk}(u, v) - \widetilde{\Sigma}_{jk}(u, v) \right| \\
 & \leq \left\| \mathbb{E}\{\widehat{\Xi}_{jk}(u, v)\} \right\|_{\min}^{-1} \left\| \widehat{\mathbf{Z}}_{jk}(u, v) - \mathbb{E}\{\widehat{\mathbf{Z}}_{jk}(u, v)\} \right\| \\
 & \quad + \left\| \mathbb{E}\{\widehat{\Xi}_{jk}(u, v)\} \right\|_{\min}^{-1} \left\| \widehat{\Xi}_{jk}(u, v) \right\|_{\min}^{-1} \left\| \widehat{\mathbf{Z}}_{jk}(u, v) \right\| \left\| \widehat{\Xi}_{jk}(u, v) - \mathbb{E}\{\widehat{\Xi}_{jk}(u, v)\} \right\|_{\text{F}}.
 \end{aligned} \tag{A.20}$$

In the following, we will prove the concentration results for case  $j \neq k$ , and the results for the case  $j = k$  can be proved in a similar manner.

### A.2.2 LOCAL CONCENTRATION INEQUALITY

We will firstly show that there exists some positive constant  $c$  (independent of  $u, v$ ) such that for any  $\delta > 0$  and  $(u, v) \in \mathcal{U}^2$ ,

$$\mathbb{P}\left\{ \left\| \widehat{\Xi}_{jk}(u, v) - \mathbb{E}\{\widehat{\Xi}_{jk}(u, v)\} \right\|_{\text{F}} \geq \delta \right\} \leq 18 \exp\left(-\frac{c\nu_{n,T,h,jk}\delta^2}{1+\delta}\right). \tag{A.21}$$

For  $m, l = 1, 2, 3$ , let  $\widehat{\Xi}_{jkm}(u, v)$  be the  $(m, l)$ th entry of  $\widehat{\Xi}_{jk}(u, v)$ . It follows from Assumptions 4 and 6 that for any integer  $q = 2, 3, \dots$  and  $s, s' = 0, 1, 2$ ,

$$\begin{aligned}
 & \mathbb{E}\left\{ \left| \left( \frac{U_{ijt} - u}{h} \right)^s \left( \frac{U_{ikt'} - v}{h} \right)^{s'} K_h(U_{ijt} - u) K_h(U_{ikt'} - v) \right|^q \right\} \\
 & \leq \int h^{-2q} K^q\left(\frac{t-u}{h}\right) K^q\left(\frac{t'-v}{h}\right) \left| \frac{t-u}{h} \right|^{sq} \left| \frac{t'-v}{h} \right|^{s'q} f_U(t) f_U(t') dt dt' \leq ch^{2-2q}.
 \end{aligned} \tag{A.22}$$

Note that Assumption 3 implies that the weights are of the same order  $w_{ijk} \asymp (n\bar{T}_{\Sigma,jk}^2)^{-1}$ . By (A.22),

$$\begin{aligned}
 & \sum_{i=1}^n \sum_{t=1}^{T_{ij}} \sum_{t'=1}^{T_{ik}} \mathbb{E}\left\{ \left| \left( \frac{U_{ijt} - u}{h} \right)^s \left( \frac{U_{ikt'} - v}{h} \right)^{s'} K_h(U_{ijt} - u) K_h(U_{ikt'} - v) \right|^2 \right\} \leq cn\bar{T}_{\Sigma,jk}^2 h^{-2}, \\
 & \sum_{i=1}^n \sum_{t=1}^{T_{ij}} \sum_{t'=1}^{T_{ik}} \mathbb{E}\left\{ \left| \left( \frac{U_{ijt} - u}{h} \right)^s \left( \frac{U_{ikt'} - v}{h} \right)^{s'} K_h(U_{ijt} - u) K_h(U_{ikt'} - v) \right|^q \right\} \leq 2^{-1} q! cn\bar{T}_{\Sigma,jk}^2 h^{2-2q}
 \end{aligned}$$

for  $q \geq 3$ . Applying the Bernstein inequality yields that there exists some positive constant  $c$  (independent of  $u, v$ ) such that for any  $\delta > 0$  and  $(u, v) \in \mathcal{U}^2$ ,

$$\mathbb{P}\left\{ \left| \widehat{\Xi}_{jkm}(u, v) - \mathbb{E}\{\widehat{\Xi}_{jkm}(u, v)\} \right| \geq \delta \right\} \leq 2 \exp\left(-\frac{c\nu_{n,T,h,jk}\delta^2}{1+\delta}\right),$$

for  $m, l = 1, 2, 3$ , which, by the union bound of probability, implies that (A.21) holds.

For  $m = 1, 2, 3$ , let  $\widehat{Z}_{jkm}(u, v)$  be the  $m$ th element of  $\widehat{\mathbf{Z}}_{jk}(u, v)$ . We will next show that, there exists some positive constant  $c$  (independent of  $u, v$ ) such that for any  $\delta > 0$  and  $(u, v) \in \mathcal{U}^2$ ,

$$\mathbb{P}\left\{ \left| \widehat{Z}_{jkm}(u, v) - \mathbb{E}\{\widehat{Z}_{jkm}(u, v)\} \right| \geq \delta \right\} \leq c \exp\left(-\frac{c\nu_{n,T,h,jk}\delta^2}{1+\delta}\right), \tag{A.23}$$

where  $\nu_{n,T,h,jk} = n(1 \wedge \bar{T}_{\Sigma,jk}^2 h^2)$ . We only need to consider the case  $m = 1$ , while the results for cases  $m = 2, 3$  can be proved similarly. Denote that

$$\begin{aligned}\zeta_{ijkts} &= \{Y_{ijt} - \mu_j(U_{ijt})\}\{Y_{iks} - \mu_k(U_{iks})\} - \Sigma_{jk}(U_{ijt}, U_{iks}), \\ \hat{Z}_{jk4}(u, v) &= \sum_{i=1}^n w_{ijk} \sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} K_h(U_{ijt} - u) K_h(U_{iks} - v) \Sigma_{jk}(U_{ijt}, U_{iks}), \\ \hat{Z}_{jk5}(u, v) &= \sum_{i=1}^n w_{ijk} \sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} K_h(U_{ijt} - u) K_h(U_{iks} - v) \zeta_{ijkts}.\end{aligned}$$

Then we rewrite  $\hat{Z}_{jk1}(u, v) - \mathbb{E}\{\hat{Z}_{jk1}(u, v)\}$  as

$$\hat{Z}_{jk1}(u, v) - \mathbb{E}\{\hat{Z}_{jk1}(u, v)\} = \hat{Z}_{jk4}(u, v) - \mathbb{E}\{\hat{Z}_{jk4}(u, v)\} + \hat{Z}_{jk5}(u, v). \quad (\text{A.24})$$

Following the same procedure to prove (A.21) with the aid of the Bernstein inequality, we can obtain that there exists some positive constant  $c$  such that for any  $\delta > 0$  and  $(u, v) \in \mathcal{U}^2$ ,

$$\mathbb{P}\left\{|\hat{Z}_{jk4}(u, v) - \mathbb{E}\{\hat{Z}_{jk4}(u, v)\}| \geq \delta\right\} \leq 2 \exp\left(-\frac{c\nu_{n,T,h,jk}\delta^2}{1+\delta}\right). \quad (\text{A.25})$$

Now we consider the tail behavior of  $\hat{Z}_{jk5}(u, v)$ . Define the event  $\tilde{V}_{jk} = \{(U_{ijt}, U_{iks}), t \in [T_{ij}], s \in [T_{ik}], i \in [n]\}$ . Note a random variable  $X$  is sub-exponential if there exist positive constants  $c_1$  and  $c_2$  such that  $\mathbb{E}(\exp[\lambda\{X - \mathbb{E}(X)\}]) \leq \exp(c_1^2\lambda^2/2)$  for all  $|\lambda| < c_2^{-1}$ . The sub-Gaussianities under Assumption 1 implies that, conditional on the event  $\tilde{V}_{jk}$ ,  $Y_{ijt} - \mu_j(U_{ijt})$  and  $Y_{iks} - \mu_k(U_{iks})$  are sub-Gaussian random variables, then  $\{Y_{ijt} - \mu_j(U_{ijt})\}\{Y_{iks} - \mu_k(U_{iks})\}$  is a sub-exponential random variable, and hence we can obtain the Bernstein-type bound  $\mathbb{E}\{\exp(\lambda\zeta_{ijkts})|\tilde{V}_{jk}\} \leq \exp\{(1-c\lambda)^{-1}c\lambda^2\}$  for any  $\lambda \in (0, c^{-1})$ . Rewrite  $\hat{Z}_{jk5}(u, v) = \sum_{i=1}^n w_{ijk}\phi_{ijk1}(u, v)$  with  $\phi_{ijk1}(u, v) = \sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} K_h(U_{ijt} - u) K_h(U_{iks} - v) \zeta_{ijkts}$ . Note that, for each  $i$ , Assumption 3 implies that  $w_{ijk} \asymp (n\bar{T}_{\Sigma,jk}^2)^{-1}$ . If  $\sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} K_h(U_{ijt} - u) K_h(U_{iks} - v) > 0$  holds, it follows from Jensen's inequality and the above result that

$$\begin{aligned}& \mathbb{E}\left[\exp\{\lambda w_{ijk}\phi_{ijk1}(u, v)\}|\tilde{V}_{jk}\right] \\ & \leq \frac{1}{\sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} K_h(U_{ijt} - u) K_h(U_{iks} - v)} \sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} \left(K_h(U_{ijt} - u) K_h(U_{iks} - v)\right) \\ & \quad \times \mathbb{E}\left[\exp\left\{\lambda w_{ijk}\zeta_{ijkts} \sum_{t'=1}^{T_{ij}} \sum_{s'=1}^{T_{ik}} K_h(U_{ijt'} - u) K_h(U_{iks'} - v)\right\}|\tilde{V}_{jk}\right] \\ & \leq \exp\left[\frac{c\lambda^2(n\bar{T}_{\Sigma,jk}^2)^{-2} \left\{\sum_{t,s} K_h(U_{ijt} - u) K_h(U_{iks} - v)\right\}^2}{1 - c\lambda(n\bar{T}_{\Sigma,jk}^2)^{-1} \sum_{t,s} K_h(U_{ijt} - u) K_h(U_{iks} - v)}\right].\end{aligned}$$

where  $0 < \lambda(n\bar{T}_{\Sigma,jk}^2)^{-1} \sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} K_h(U_{ijt} - u) K_h(U_{iks} - v) < c^{-1}$ . It is obvious that the above inequality still holds even if  $\sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} K_h(U_{ijt} - u) K_h(U_{iks} - v) = 0$ . Assumption 5

implies that the number of nonzero terms in  $\sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} K_h(U_{ijt} - u)K_h(U_{iks} - v)$  has an upper bound  $c(1 \vee \bar{T}_{\Sigma, jk}^2 h^2)$ , which yields that  $\sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} K_h(U_{ijt} - u)K_h(U_{iks} - v) \leq ch^{-2}(1 \vee \bar{T}_{\Sigma, jk}^2 h^2)$ . Therefore, for any  $\lambda$  satisfying  $0 < \lambda(n\bar{T}_{\Sigma, jk}^2 h^2)^{-1}(1 \vee \bar{T}_{\Sigma, jk}^2 h^2) < c^{-1}$  for some constant  $c > 0$ , we obtain that

$$\begin{aligned} & \mathbb{E} \left[ \exp \left\{ \lambda \sum_{i=1}^n w_{ijk} \phi_{ijk1}(u, v) \right\} \middle| \tilde{V}_{jk} \right] \\ & \leq \exp \left\{ \frac{c\lambda^2(n\bar{T}_{\Sigma, jk}^2)^{-2}h^{-2}(1 \vee \bar{T}_{\Sigma, jk}^2 h^2) \sum_{i=1}^n \sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} K_h(U_{ijt} - u)K_h(U_{iks} - v)}{1 - c\lambda(n\bar{T}_{\Sigma, jk}^2 h^2)^{-1}(1 \vee \bar{T}_{\Sigma, jk}^2 h^2)} \right\}. \end{aligned}$$

For any  $\delta > 0$ , define the event

$$\Lambda_{jk,1}(\delta) = \left\{ \sum_{i=1}^n \sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} K_h(U_{ijt} - u)K_h(U_{iks} - v) \leq c(1 + \delta)n\bar{T}_{\Sigma, jk}^2 \right\}.$$

We have

$$\mathbb{E} \left[ \exp \left\{ \lambda \sum_{i=1}^n w_{ijk} \phi_{ijk1}(u, v) \right\} \middle| \Lambda_{jk,1}(\delta) \right] \leq \exp \left\{ \frac{c\lambda^2(1 + \delta)(n\bar{T}_{\Sigma, jk}^2 h^2)^{-1}(1 \vee \bar{T}_{\Sigma, jk}^2 h^2)}{1 - c\lambda(n\bar{T}_{\Sigma, jk}^2 h^2)^{-1}(1 \vee \bar{T}_{\Sigma, jk}^2 h^2)} \right\}.$$

Consequently, we obtain that

$$\mathbb{P} \left\{ \sum_{i=1}^n w_{ijk} \phi_{ijk1}(u) \geq \delta \middle| \Lambda_{jk,1}(\delta) \right\} \leq \exp \left\{ -\lambda\delta + \frac{c\lambda^2(1 + \delta)(n\bar{T}_{\Sigma, jk}^2 h^2)^{-1}(1 \vee \bar{T}_{\Sigma, jk}^2 h^2)}{1 - c\lambda(n\bar{T}_{\Sigma, jk}^2 h^2)^{-1}(1 \vee \bar{T}_{\Sigma, jk}^2 h^2)} \right\}. \quad (\text{A.26})$$

With the choice of  $\lambda = n\bar{T}_{\Sigma, jk}^2 h^2 \delta \{2c(1 + \delta)(1 \vee \bar{T}_{\Sigma, jk}^2 h^2) + c\delta(1 \vee \bar{T}_{\Sigma, jk}^2 h^2)\}^{-1}$ , (A.26) reduces to

$$\mathbb{P} \left\{ \sum_{i=1}^n w_{ijk} \phi_{ijk1}(u, v) \geq \delta \middle| \Lambda_{jk,1}(\delta) \right\} \leq \exp \left\{ -\frac{c\nu_{n,T,h,jk}\delta^2}{1 + \delta} \right\}, \quad (\text{A.27})$$

where the constant  $c$  is chosen to satisfy  $c\lambda(n\bar{T}_{\Sigma, jk}^2 h^2)^{-1}(1 \vee \bar{T}_{\Sigma, jk}^2 h^2) \leq 1/2$ . Note that

$$\sum_{i=1}^n \sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} \mathbb{E} \{ K_h(U_{ijt} - u)K_h(U_{iks} - v) \} \leq cn\bar{T}_{\Sigma, jk}^2.$$

By the Bernstein inequality, we obtain that there exists some positive constant  $c$  such that for any  $\delta > 0$

$$\begin{aligned} & \mathbb{P} \left( \sum_{i=1}^n \sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} \left[ K_h(U_{ijt} - u)K_h(U_{iks} - v) - \mathbb{E} \{ K_h(U_{ijt} - u)K_h(U_{iks} - v) \} \right] \geq n\bar{T}_{\Sigma, jk}^2 \delta \right) \\ & \leq \exp \left( \frac{-c\nu_{n,T,h,jk}\delta^2}{1 + \delta} \right), \end{aligned}$$

which implies that

$$1 - \mathbb{P}\{\Lambda_{jk,1}(\delta)\} \leq \exp\left(-\frac{c\nu_{n,T,h,jk}\delta^2}{1+\delta}\right). \quad (\text{A.28})$$

Combining (A.27) and (A.28), we obtain that there exists some constant  $c > 0$  such that for any  $\delta > 0$ ,

$$\mathbb{P}\left\{\widehat{Z}_{jk5}(u, v) \geq \delta\right\} \leq \mathbb{P}\left\{\widehat{Z}_{jk5}(u, v) \geq \delta \mid \Lambda_{jk,1}(\delta)\right\} + \mathbb{P}\left\{\Lambda_{jk,1}(\delta)^c\right\} \leq 2 \exp\left(-\frac{c\nu_{n,T,h,jk}\delta^2}{1+\delta}\right),$$

which leads to

$$\mathbb{P}\left\{|\widehat{Z}_{jk5}(u, v)| \geq \delta\right\} \leq 4 \exp\left(-\frac{c\nu_{n,T,h,jk}\delta^2}{1+\delta}\right).$$

It follows from the above, (A.24) and (A.25) that for each  $\delta > 0$  and  $(u, v) \in \mathcal{U}^2$ , there exists some positive constant  $c$  such that

$$\mathbb{P}\left\{|\widehat{Z}_{jk1}(u, v) - \mathbb{E}\{\widehat{Z}_{jk1}(u, v)\}| \geq \delta\right\} \leq 6 \exp\left(-\frac{c\nu_{n,T,h,jk}\delta^2}{1+\delta}\right).$$

Define the event  $\Lambda_{jk,2}(\delta) = \{\|\widehat{\Xi}_{jk}(u, v) - \mathbb{E}\{\widehat{\Xi}_{jk}(u, v)\}\|_{\text{F}} \leq \delta/2\}$ . Note that  $\mathbb{E}\{\widehat{\Xi}_{jk}(u, v)\}$  is positive definite. On the event  $\Lambda_{jk,2}(\delta)$  with  $\delta \in (0, 1]$ , we obtain that

$$\|\widehat{\Xi}_{jk}(u, v)\|_{\min} \geq c(1 - \delta/2). \quad (\text{A.29})$$

By (A.21), we have

$$1 - \mathbb{P}\{\Lambda_{jk,2}(\delta)\} \leq 18 \exp\left(-\frac{c\nu_{n,T,h,jk}\delta^2}{1+\delta}\right). \quad (\text{A.30})$$

Define the event  $\Lambda_{jk,3}(\delta) = \{\|\widehat{\mathbf{Z}}_{jk}(u, v) - \mathbb{E}\{\widehat{\mathbf{Z}}_{jk}(u, v)\}\| \leq \delta\}$ . Note that, under Assumption 3 with  $w_{ijk} \asymp (n\bar{T}_{\Sigma,jk}^2)^{-1}$ ,  $\sum_{i=1}^n w_{ijk} \sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} \mathbb{E}\{K_h(U_{ijt} - u)K_h(U_{iks} - v)\} \leq c$ , hence  $\|\mathbb{E}\{\widehat{\mathbf{Z}}_{jk}(u, v)\}\|$  is uniformly bounded over  $\mathcal{U}^2$ . On the event  $\Omega_{jk,3}(\delta)$ , we have

$$\|\widehat{\mathbf{Z}}_{jk}(u, v)\| \leq c(1 + \delta). \quad (\text{A.31})$$

On the event  $\Lambda_{jk,2}(\delta) \cap \Lambda_{jk,3}(\delta)$  with  $\delta \in (0, 1]$ , it follows from (A.20), (A.29) and (A.31) that

$$|\widehat{\Sigma}_{jk}(u, v) - \widetilde{\Sigma}_{jk}(u, v)| \leq c\delta + c(1 - \delta/2)^{-1}(1 + \delta)\delta \leq c_3\delta.$$

This together with concentration inequalities in (A.23) and (A.30) implies that there exist some positive universal constants  $c_1$  and  $c_2$  such that for any  $\delta \in (0, 1]$  and  $(u, v) \in \mathcal{U}^2$ ,

$$\mathbb{P}\left\{|\widehat{\Sigma}_{jk}(u, v) - \widetilde{\Sigma}_{jk}(u, v)| \geq \delta\right\} \leq c_2 \exp(-c_1\nu_{n,T,h,jk}\delta^2),$$

which completes the proof of local concentration inequality for the covariance estimator.

## A.2.3 CONCENTRATION INEQUALITY IN HILBERT–SCHMIDT NORM

We will derive the  $L_2$  concentration inequality of  $\|\widehat{\Sigma}_{jk} - \widetilde{\Sigma}_{jk}\|_S$ . Let

$$\widetilde{\Xi}_{jk}(u, v) = (n\bar{T}_{\Sigma, jk}^2)^{-1} \sum_{i=1}^n \sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} \widetilde{U}_{ij kts} \widetilde{U}_{ij kts}^T K_h(U_{ijt} - u) K_h(U_{iks} - v).$$

Then we have that  $\|\widehat{\Xi}_{jk}(u, v)\|_{\min} \geq c \|\widetilde{\Xi}_{jk}(u, v)\|_{\min}$ . Similar to Appendix A.1.3, we will give a lower bound on  $\|\widetilde{\Xi}_{jk}(u, v)\|_{\min}$ . Denote  $\widetilde{W} = \sup_{(u,v) \in \mathcal{U}^2} \|\widetilde{\Xi}_{jk}(u, v) - \mathbb{E}\{\widetilde{\Xi}_{jk}(u, v)\}\|_F$ . For  $t, s = 1, 2, 3$ , let  $\widetilde{\Xi}_{j kts}(u, v)$  be the  $(t, s)$ th entry of  $\widetilde{\Xi}_{jk}(u, v)$ . Note that  $\mathbb{E}\{|(U_{ijt} - u)^a h^{-a} (U_{iks} - v)^b h^{-b} K_h(U_{ijt} - u) K_h(U_{iks} - v)|\} \leq c$  for  $a, b = 0, 1, 2$ , and, moreover,  $\mathbb{E}(\widetilde{W}) \leq 6 \max_{t,s} \mathbb{E}\{\sup_{(u,v) \in \mathcal{U}^2} |\widetilde{\Xi}_{j kts}(u, v)|\}$ . In an analogy to Lemma 13.5 of Boucheron et al. (2013) and by the similar arguments below (A.14) in Appendix A.1.3, we can show that  $\mathbb{E}(\widetilde{W}) \leq c(n\bar{T}_{\Sigma, jk}^2)^{-1/2}$ . In addition, it follows from the facts  $\text{var}(W) \leq \mathbb{E}(W^2) \leq 9 \max_{t,s} \text{var}\{\sup_{(u,v) \in \mathcal{U}^2} \widetilde{\Xi}_{j kts}(u, v)\}$ ,  $|(U_{ijt} - u)^a h^{-a} (U_{iks} - v)^b h^{-b} K_h(U_{ijt} - u) K_h(U_{iks} - v)| \leq ch^{-2}$ ,  $\mathbb{E}\{(U_{ijt} - u)^{2a} h^{-2a} (U_{iks} - v)^{2b} h^{-2b} K_h^2(U_{ijt} - u) K_h^2(U_{iks} - v)\} \leq ch^{-2}$  for  $a, b = 0, 1, 2$ , and Theorem 11.10 of Boucheron et al. (2013) that  $\text{var}(n\bar{T}_{\Sigma, jk}^2 h^2 \widetilde{W}) \leq 2\mathbb{E}(n\bar{T}_{\Sigma, jk}^2 h^2 \widetilde{W}) + \sum_{i=1}^n \sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} ch^{-2} h^4 \leq c(n\bar{T}_{\Sigma, jk}^2)^{1/2} h^2 + cn\bar{T}_{\Sigma, jk}^2 h^2$ , which implies that the variance of  $\widetilde{W}$  is bounded by  $c(n\bar{T}_{\Sigma, jk}^2 h^2)^{-1} \leq c(\nu_{n,T,h,jk})^{-1}$ . Noting the similar arguments below (A.14) and applying Theorem 12.5 of Boucheron et al. (2013), we obtain that there exists some positive constant  $c$  such that, for any  $\delta > 0$ ,

$$\mathbb{P}\left\{\widetilde{W} - \mathbb{E}(\widetilde{W}) > \delta\right\} \leq \exp\left(-\frac{c\nu_{n,T,h,jk}\delta^2}{1+\delta}\right). \quad (\text{A.32})$$

Define the event  $\Lambda_{jk,4}(\delta) = \{\sup_{(u,v) \in \mathcal{U}^2} \|\widetilde{\Xi}_{jk}(u, v) - \mathbb{E}\{\widetilde{\Xi}_{jk}(u, v)\}\|_F \leq \delta/2\}$  with  $\delta \in (0, 1]$ . By (A.32), we obtain that there exists some constant  $c > 0$  such that, for any  $\delta \in (0, 1]$ ,

$$1 - \mathbb{P}\{\Lambda_{jk,4}(\delta)\} \leq 2 \exp(-c\nu_{n,T,h,jk}\delta^2).$$

On the event  $\Lambda_{jk,4} = \Lambda_{jk,4}(\tilde{\delta}_1)$  with  $c(\nu_{n,T,h,jk})^{-1/2} < \tilde{\delta}_1 \leq 1$ , we have  $\|\widehat{\Xi}_{jk}(u, v)\|_{\min} \geq c \|\widetilde{\Xi}_{jk}(u, v)\|_{\min} \geq c(1 - \tilde{\delta}_1/2) \geq c/2$ . Notice that  $\mathbb{E}\{\widehat{\Xi}_{jk}(u, v)\}$  is positive definite and  $\|\mathbb{E}\{\widehat{\mathbf{Z}}_{jk}(u, v)\}\|$  is uniformly bounded over  $\mathcal{U}^2$ . On the event  $\Lambda_{jk,4}$ , it thus follows from (A.20) and  $\|\widehat{\mathbf{Z}}_{jk}(u, v)\| \leq \|\widehat{\mathbf{Z}}_{jk}(u, v) - \mathbb{E}\{\widehat{\mathbf{Z}}_{jk}(u, v)\}\| + \|\mathbb{E}\{\widehat{\mathbf{Z}}_{jk}(u, v)\}\|$  that

$$|\widehat{\Sigma}_{jk}(u, v) - \widetilde{\Sigma}_{jk}(u, v)| \leq c \|\widehat{\mathbf{Z}}_{jk}(u, v) - \mathbb{E}\{\widehat{\mathbf{Z}}_{jk}(u, v)\}\| + c \|\mathbb{E}\{\widehat{\Xi}_{jk}(u, v) - \mathbb{E}\{\widehat{\Xi}_{jk}(u, v)\}\|_F \quad (\text{A.33})$$

and the positive constant  $c$  does not depend on  $(u, v) \in \mathcal{U}^2$ .

Combining (A.33) with (A.21), (A.23) and applying the first part of Lemma 11 yields that, for any  $(u, v) \in \mathcal{U}^2$  and integer  $q \geq 1$ ,

$$\mathbb{E}\left\{|\widehat{\Sigma}_{jk}(u, v) - \widetilde{\Sigma}_{jk}(u, v)|^{2q} \mid \Lambda_{jk,4}\right\} \leq q! c \left(\frac{4}{c\nu_{n,T,h,jk}}\right)^q + (2q)! c \left(\frac{4}{c\nu_{n,T,h,jk}}\right)^{2q}.$$

Applying the second part of Lemma 11, we can show that, for each  $\delta \in (0, 1]$ ,

$$\mathbb{P}\left(\|\widehat{\Sigma}_{jk} - \widetilde{\Sigma}_{jk}\|_S \geq \delta\right) \leq \mathbb{P}\left(\|\widehat{\Sigma}_{jk} - \widetilde{\Sigma}_{jk}\|_S \geq \delta \mid \Lambda_{jk,4}\right) + \mathbb{P}(\Lambda_{jk,4}^c) \leq c_4 \exp(-c_3 \nu_{n,T,h,jk} \delta^2),$$

which means that (7) in Theorem 3 holds and completes the proof of concentration inequality for the covariance estimator in Hilbert–Schmidt norm.

## A.2.4 CONCENTRATION INEQUALITY IN SUPREMUM NORM

We will derive the uniform concentration bound of  $\sup_{(u,v) \in \mathcal{U}^2} |\widehat{\Sigma}_{jk}(u, v) - \widetilde{\Sigma}_{jk}(u, v)|$ . We partition the interval  $\mathcal{U} = [0, 1]$  into  $N$  subintervals  $I_s$  for  $s \in [N]$  of equal length. Let  $u_s$  and  $v_{s'}$  be the centers of  $I_s$  and  $I_{s'}$ , respectively, then we have

$$\begin{aligned} \sup_{(u,v) \in \mathcal{U}^2} |\widehat{\Sigma}_{jk}(u, v) - \widetilde{\Sigma}_{jk}(u, v)| &\leq \max_{s, s' \in [N]} \left[ \left| \widehat{\Sigma}_{jk}(u_s, v_{s'}) - \widetilde{\Sigma}_{jk}(u_s, v_{s'}) \right| \right. \\ &\quad \left. + \left| \{\widehat{\Sigma}_{jk}(u, v) - \widehat{\Sigma}_{jk}(u_s, v_{s'})\} - \{\widetilde{\Sigma}_{jk}(u, v) - \widetilde{\Sigma}_{jk}(u_s, v_{s'})\} \right| \right]. \end{aligned}$$

We need to bound the second term. By some calculations, it suffices to bound  $\left| \{\widehat{Z}_{jkm}(u, v) - \widehat{Z}_{jkm}(u_s, v_{s'})\} - [\mathbb{E}\{\widehat{Z}_{jkm}(u, v)\} - \mathbb{E}\{\widehat{Z}_{jkm}(u_s, v_{s'})\}] \right|$  and  $\left| \{\widehat{\Xi}_{jkl}(u, v) - \widehat{\Xi}_{jkl}(u_s, v_{s'})\} - [\mathbb{E}\{\widehat{\Xi}_{jkl}(u, v)\} - \mathbb{E}\{\widehat{\Xi}_{jkl}(u_s, v_{s'})\}] \right|$  for  $m, l = 1, 2, 3$ , which means that we need to bound  $|\widehat{Z}_{jkm}(u, v) - \widehat{Z}_{jkm}(u_s, v_{s'})|$  and  $|\widehat{\Xi}_{jkl}(u, v) - \widehat{\Xi}_{jkl}(u_s, v_{s'})|$ . Let  $(u, v) \in I_s \times I_{s'}$  and consider  $|\widehat{Z}_{jk1}(u, v) - \widehat{Z}_{jk1}(u_s, v_{s'})|$  for the case of  $j \neq k$  first. The results for the case of  $j = k$  can be proved in a similar fashion. Define the event  $\Lambda_{Z,jk1} = \left\{ \sum_{i=1}^n w_{ijk} \sum_{t=1}^{T_{ij}} \sum_{t'=1}^{T_{ik}} |\Theta_{ijkt't'}| \leq \mathbb{E}(\sum_{i=1}^n w_{ijk} \sum_{t=1}^{T_{ij}} \sum_{t'=1}^{T_{ik}} |\Theta_{ijkt't'}|) + 1 \right\}$ . On this event, it follows from Assumption 6(ii) that

$$\begin{aligned} &\left| \widehat{Z}_{jk1}(u, v) - \widehat{Z}_{jk1}(u_s, v_{s'}) \right| \\ &\leq \left| \sum_{i=1}^n w_{ijk} \sum_{t=1}^{T_{ij}} \sum_{t'=1}^{T_{ik}} \Theta_{ijkt't'} \left[ \{K_h(U_{ijt} - u) - K_h(U_{ijt} - u_s)\} K_h(U_{ikt'} - v) \right. \right. \\ &\quad \left. \left. + \{K_h(U_{ikt'} - v) - K_h(U_{ikt'} - v_{s'})\} K_h(U_{ijt} - u_s) \right] \right| \\ &\leq \frac{c(|u - u_s| \vee |v - v_{s'}|)}{h^2} \sum_{i=1}^n w_{ijk} \sum_{t=1}^{T_{ij}} \sum_{t'=1}^{T_{ik}} |\Theta_{ijkt't'}| \{K_h(U_{ikt'} - v) + K_h(U_{ijt} - u_s)\} \\ &\leq \frac{c}{Nh^3} \left\{ \mathbb{E} \left( \sum_{i=1}^n w_{ijk} \sum_{t=1}^{T_{ij}} \sum_{t'=1}^{T_{ik}} |\Theta_{ijkt't'}| \right) + 1 \right\} \leq \frac{c}{Nh^3}. \end{aligned}$$

Applying similar techniques as above, we can define events  $\Lambda_{Z,jkm}$  and  $\Lambda_{\Xi,jkl}$  for  $m, l = 1, 2, 3$ . On the intersection of these events, we can obtain that  $|\widehat{Z}_{jkm}(u, v) - \widehat{Z}_{jkm}(u_s, v_{s'})| \leq c(Nh^3)^{-1}$  and  $|\widehat{\Xi}_{jkl}(u, v) - \widehat{\Xi}_{jkl}(u_s, v_{s'})| \leq c(Nh^3)^{-1}$ . Combing the above results, we have

$$\sup_{(u,v) \in \mathcal{U}^2} |\widehat{\Sigma}_{jk}(u, v) - \widetilde{\Sigma}_{jk}(u, v)| \leq \max_{s, s' \in [N]} \left| \widehat{\Sigma}_{jk}(u_s, v_{s'}) - \widetilde{\Sigma}_{jk}(u_s, v_{s'}) \right| + \frac{c}{Nh^3}.$$

By the Bernstein inequality, we have  $\mathbb{P}(\Lambda_{Z,jkm}^c) \leq \exp\{-\lambda + c_1^2 \lambda^2 (n - c_2 \lambda)^{-1}\}$  for  $\lambda \in (0, nc^{-1}]$ . With  $\lambda = n(2c_1^2 + c_2)^{-1}$ , the right-side reduces to  $\exp(-cn) \leq \exp(-c\nu_{n,T,h,jk})$  for  $m, l = 1, 2, 3$ . Similarly,  $\mathbb{P}(\Lambda_{\Xi,jkl}^c) \leq \exp(-cn) \leq \exp(-c\nu_{n,T,h,jk})$  for  $m, l = 1, 2, 3$ . It follows from the above results and the union bound of probability with the choice of

$N = \lfloor c(h^3\delta)^{-1} \rfloor$  that there exist some positive constants  $c_3$  and  $c_4$  such that, for any  $\delta \in (0, 1]$ ,

$$\mathbb{P}\left\{ \sup_{(u,v) \in \mathcal{U}^2} |\widehat{\Sigma}_{jk}(u,v) - \widetilde{\Sigma}_{jk}(u,v)| \geq \delta \right\} \leq \frac{c_4}{h^6\delta^2} \exp(-c_3\nu_{n,T,h,jk}\delta^2). \quad (\text{A.34})$$

Take arbitrarily small  $\epsilon_2 > 0$ . If  $n^{\epsilon_2}\nu_{n,T,h,jk}\delta^2 \geq 1$ , the right side of (A.34) reduces to  $c_4n^{\epsilon_2}\nu_{n,T,h,jk}h^{-6} \exp(-c_3\nu_{n,T,h,jk}\delta^2)$ . If  $n^{\epsilon_2}\nu_{n,T,h,jk}\delta^2 \leq 1$ , we can choose  $c_4$  and  $n^{\epsilon_2} > c$  such that  $c_4 \exp(-c_3c^{-1}) \geq 1$  and hence the same bound  $c_4n^{\epsilon_2}\nu_{n,T,h,jk}h^{-6} \exp(-c_3\nu_{n,T,h,jk}\delta^2)$  can still be used. We complete the proof of (8) in Theorem 3, the concentration inequality for the covariance estimator in supremum norm.  $\blacksquare$

### A.3 Proof of Theorem 6

Note that  $\|\widehat{\mu}_j - \mu_j\|_2 \leq \|\widehat{\mu}_j - \widetilde{\mu}_j\|_2 + \|\widetilde{\mu}_j - \mu_j\|_2$ , it suffices to bound  $\|\widetilde{\mu}_j - \mu_j\|_2$ . By (A.1), for any  $u \in \mathcal{U}$ ,

$$\widetilde{\mu}_j(u) - \mu_j(u) = \mathbf{e}_0^T \left[ \mathbb{E}\{\widehat{\mathbf{S}}_j(u)\} \right]^{-1} \mathbb{E}\left[ \widehat{\mathbf{R}}_j(u) - \widehat{\mathbf{S}}_j(u)\{\mu_j(u), 0\}^T \right].$$

By the Taylor expansion, we have

$$\mathbb{E}_\varepsilon \left[ \widehat{\mathbf{R}}_j(u) - \widehat{\mathbf{S}}_j(u)\{\mu_j(u), 0\}^T \middle| V_j \right] = \sum_{i=1}^n v_{ij} \sum_{t=1}^{T_{ij}} \widetilde{\mathbf{U}}_{ijt} K_{h_{\mu,j}}(U_{ijt} - u) \{\mu_j(U_{ijt}) - \mu_j(u)\} := \mathbf{J}_1 + \mathbf{J}_2,$$

with

$$\begin{aligned} \mathbf{J}_1 &= \sum_{i=1}^n v_{ij} \sum_{t=1}^{T_{ij}} \widetilde{\mathbf{U}}_{ijt} K_{h_{\mu,j}}(U_{ijt} - u) \frac{U_{ijt} - u}{h_{\mu,j}} h_{\mu,j} \frac{\partial \mu_j(u)}{\partial u}, \\ \mathbf{J}_2 &= \frac{1}{2} \sum_{i=1}^n v_{ij} \sum_{t=1}^{T_{ij}} \widetilde{\mathbf{U}}_{ijt} K_{h_{\mu,j}}(U_{ijt} - u) \left( \frac{U_{ijt} - u}{h_{\mu,j}} \right)^2 h_{\mu,j}^2 \frac{\partial \mu_j^2(\delta_{ijt})}{\partial u^2}, \end{aligned}$$

where  $\delta_{ijt} \in [u - h_{\mu,j}, u + h_{\mu,j}]$  and  $\mathbb{E}_\varepsilon$  denotes the expectation over  $\{Y_{ijt}\}$  in (4) conditional on the event  $V_j = \{U_{ijt}, t \in [T_{ij}], i \in [n]\}$ . First consider  $\mathbf{J}_1$ , which equals to the second column of  $\widehat{\mathbf{S}}_j(u)$  multiplied by  $h_{\mu,j} \partial \mu_j(u) / \partial u$ , hence

$$\mathbb{E}_U \left[ \mathbf{e}_0^T \left[ \mathbb{E}\{\widehat{\mathbf{S}}_j(u)\} \right]^{-1} \mathbf{J}_1 \right] = h_{\mu,j} \frac{\partial \mu_j(u)}{\partial u} \mathbf{e}_0^T \left[ \mathbb{E}\{\widehat{\mathbf{S}}_j(u)\} \right]^{-1} \mathbb{E}\{\widehat{\mathbf{S}}_j(u)\} (0, 1)^T = 0,$$

where  $\mathbb{E}_U$  denotes the expectation over  $V_j$ . Consider  $\mathbf{J}_2$  next. Under Assumption 7, we have  $K_1 = \sup_{j \in [p], \xi \in \mathcal{U}} |\partial \mu_j^2(\xi) / \partial u^2| < \infty$ . Each entry of  $|\mathbf{J}_2|$  is bounded by the (1, 1)th entry of  $\widehat{\mathbf{S}}_j(u)$  multiplied by  $K_1 h_{\mu,j}^2 / 2$ , and by  $\mathbb{E}\{K_h(U_{ijt} - u)\} \leq 1$  we have that the (1, 1)th entry of  $\mathbb{E}\{\widehat{\mathbf{S}}_j(u)\}$  is bounded by 1. Note that  $\mathbb{E}\{\widehat{\mathbf{S}}_j(u)\}$  is positive definite. These results together yield that

$$|\mathbb{E}_U(\mathbf{e}_0^T \left[ \mathbb{E}\{\widehat{\mathbf{S}}_j(u)\} \right]^{-1} \mathbf{J}_2)| \leq \|\mathbb{E}\{\widehat{\mathbf{S}}_j(u)\}^{-1}\|_{\min} \|\mathbb{E}_U(|\mathbf{J}_2|)\| \leq cK_1 h_{\mu,j}^2,$$

which implies that  $|\tilde{\mu}_j(u) - \mu_j(u)| \leq cK_1 h_{\mu,j}^2$  for any  $u \in \mathcal{U}$ . Hence

$$\sup_{u \in \mathcal{U}} |\tilde{\mu}_j(u) - \mu_j(u)| = O(h_{\mu,j}^2) \quad \text{and} \quad \|\tilde{\mu}_j - \mu_j\|_2 = O(h_{\mu,j}^2). \quad (\text{A.35})$$

For  $M > 0$  with the choice of  $\delta = (\log p)^{1/2} (\min_j \gamma_{n,T,h,j})^{-1/2} M \leq 1$ , it follows from the union bound of probability and (5) in Theorem 2 that

$$\begin{aligned} & \mathbb{P} \left\{ \frac{\max_{j \in [p]} \|\hat{\mu}_j - \tilde{\mu}_j\|_2}{(\log p)^{1/2} (\min_j \gamma_{n,T,h,j})^{-1/2}} \geq M \right\} \\ & \leq \sum_{j=1}^p \mathbb{P} \left\{ \|\hat{\mu}_j - \tilde{\mu}_j\|_2 \geq M (\log p)^{1/2} (\min_j \gamma_{n,T,h,j})^{-1/2} \right\} \\ & \leq \sum_{j=1}^p c \exp \left( -c \gamma_{n,T,h,j} M^2 \frac{\log p}{\min_j \gamma_{n,T,h,j}} \right) \leq c \exp \{ (1 - cM^2) \log p \}. \end{aligned} \quad (\text{A.36})$$

We can choose a large  $M$  such that  $1 - cM^2 < 0$ , the right-side of (A.36) tends to 0. Hence  $\max_{j \in [p]} \|\hat{\mu}_j - \tilde{\mu}_j\|_2 = O_P \{ (\log p)^{1/2} (\min_j \gamma_{n,T,h,j})^{-1/2} \}$ . Combing this with (A.35) yields that

$$\max_{j \in [p]} \|\hat{\mu}_j - \mu_j\|_2 = O_P \left\{ \left( \frac{\log p}{\min_j \gamma_{n,T,h,j}} \right)^{1/2} + \max_j h_{\mu,j}^2 \right\},$$

which completes the proof of (9).

The rate of convergence in (10) can be proved following a similar procedure. Let  $h_{\mu,\min} = \min_j h_{\mu,j}$ , we assume that  $h_{\mu,\min} \asymp \{\log(p \vee n)/n\}^{\kappa_1}$  where  $\kappa_1 \in (0, 1/2]$ . Consider  $h_{\mu,\min}$  with some  $\kappa_1^* > 1/2$ , the corresponding rate is not faster than that with  $\kappa_1 \in (0, 1/2]$ . To be specific, under the sparse design, the rate of  $\max_{j \in [p]} \sup_{u \in \mathcal{U}} |\hat{\mu}_j(u) - \mu_j(u)|$  is  $\{\log(p \vee n)/n\}^{1/2 - \kappa_1^*/2}$ , which is slower than  $\{\log(p \vee n)/n\}^{2/5}$  with  $\kappa_1 = 1/5$ . Under the dense design with  $\bar{T}_\mu \{\log(p \vee n)/n\}^{\kappa_1^*} \rightarrow 0$  and  $\bar{T}_\mu \{\log(p \vee n)/n\}^{3/2} \rightarrow 0$ , the rate is  $\{\log(p \vee n)/n\}^{1/2 - \kappa_1^*/2} \bar{T}_\mu^{-1/2}$ , which is slower than  $\{\log(p \vee n)/n\}^{2/5} \bar{T}_\mu^{-2/5}$  with  $\kappa_1 \in (1/5, 1/2]$ . Under the dense design with  $\bar{T}_\mu \{\log(p \vee n)/n\}^{\kappa_1^*} \rightarrow 0$  and  $\bar{T}_\mu \{\log(p \vee n)/n\}^{3/2} \rightarrow \tilde{c}$  or  $\infty$ , the rate is  $\{\log(p \vee n)/n\}^{1/2 - \kappa_1^*/2} \bar{T}_\mu^{-1/2}$ , which is slower than  $\{\log(p \vee n)/n\}^{1/2}$  with  $\kappa_1 = 1/4$ . Under the dense design with  $\bar{T}_\mu \{\log(p \vee n)/n\}^{\kappa_1^*} \rightarrow \tilde{c}$  or  $\infty$ , the rate is  $\{\log(p \vee n)/n\}^{1/2}$ , which is the same as  $\{\log(p \vee n)/n\}^{1/2}$  with  $\kappa_1 = 1/4$ . Based on the above four cases, if  $\kappa_1^* > 1/2$ , the corresponding rate is not faster than that with some  $\kappa_1 \in (0, 1/2]$  and hence the  $\kappa_1$  that corresponds with the optimal bandwidth under sparse or dense design is in  $(0, 1/2]$ . For  $M > 0$  with the choice of  $\delta = \{\log(p \vee n)\}^{1/2} (\min_j \gamma_{n,T,h,j})^{-1/2} M \leq 1$ , by the union bound of probability and (6) in Theorem 2, we have

$$\begin{aligned} & \mathbb{P} \left\{ \frac{\max_{j \in [p]} \sup_{u \in \mathcal{U}} |\hat{\mu}_j(u) - \tilde{\mu}_j(u)|}{\{\log(p \vee n)\}^{1/2} (\min_j \gamma_{n,T,h,j})^{-1/2}} \geq M \right\} \\ & \leq \sum_{j=1}^p \mathbb{P} \left\{ \sup_{u \in \mathcal{U}} |\hat{\mu}_j(u) - \tilde{\mu}_j(u)| \geq M \{\log(p \vee n)\}^{1/2} (\min_j \gamma_{n,T,h,j})^{-1/2} \right\} \\ & \leq \sum_{j=1}^p \frac{c(n^{\epsilon_1} \gamma_{n,T,h,j})^{1/2}}{h_{\mu,\min}^2} \exp \left\{ -c \gamma_{n,T,h,j} M^2 \frac{\log(p \vee n)}{\min_j \gamma_{n,T,h,j}} \right\} \\ & \leq c \exp \left\{ \left( \frac{3 + \epsilon_1}{2} + 2\kappa_1 - cM^2 \right) \log(p \vee n) \right\}. \end{aligned} \quad (\text{A.37})$$

We can choose a large  $M$  such that  $3 + \epsilon_1 + 4\kappa_1 - 2cM^2 < 0$ , the right-side of (A.37) tends to 0. Hence

$$\max_{j \in [p]} \sup_{u \in \mathcal{U}} |\hat{\mu}_j(u) - \tilde{\mu}_j(u)| = O_P \left[ \left\{ \frac{\log(p \vee n)}{\min_j \gamma_{n,T,h,j}} \right\}^{1/2} \right].$$

Combing this with (A.35) yields that

$$\max_{j \in [p]} \sup_{u \in \mathcal{U}} |\hat{\mu}_j(u) - \mu_j(u)| = O_P \left[ \left\{ \frac{\log(p \vee n)}{\min_j \gamma_{n,T,h,j}} \right\}^{1/2} + \max_j h_{\mu,j}^2 \right],$$

which completes the proof of (10).  $\blacksquare$

#### A.4 Proof of Theorem 7

Note  $\|\hat{\Sigma}_{jk} - \Sigma_{jk}\|_{\mathcal{S}} \leq \|\hat{\Sigma}_{jk} - \tilde{\Sigma}_{jk}\|_{\mathcal{S}} + \|\tilde{\Sigma}_{jk} - \Sigma_{jk}\|_{\mathcal{S}}$ , it suffices to bound  $\|\tilde{\Sigma}_{jk} - \Sigma_{jk}\|_{\mathcal{S}}$ . By (A.19), for any  $(u, v) \in \mathcal{U}^2$ ,

$$\tilde{\Sigma}_{jk}(u, v) - \Sigma_{jk}(u, v) = \tilde{\mathbf{e}}_0^T \left[ \mathbb{E} \{ \hat{\mathbf{\Xi}}_{jk}(u, v) \} \right]^{-1} \mathbb{E} \left[ \hat{\mathbf{Z}}_{jk}(u, v) - \hat{\mathbf{\Xi}}_{jk}(u, v) \{ \Sigma_{jk}(u, v), 0, 0 \}^T \right].$$

By the Taylor expansion, we have

$$\begin{aligned} & \mathbb{E}_\epsilon \left[ \hat{\mathbf{Z}}_{jk}(u, v) - \hat{\mathbf{\Xi}}_{jk}(u, v) \{ \Sigma_{jk}(u, v), 0, 0 \}^T \mid \tilde{V}_{jk} \right] \\ &= \sum_{i=1}^n w_{ijk} \sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} \tilde{\mathbf{U}}_{ij kts} K_{h_{\Sigma,jk}}(U_{ijt} - u) K_{h_{\Sigma,jk}}(U_{iks} - v) \{ \Sigma_{jk}(U_{ijt}, U_{iks}) - \Sigma_{jk}(u, v) \} \\ &:= \mathbf{L}_1 + \mathbf{L}_2 + \mathbf{L}_3, \end{aligned}$$

where

$$\begin{aligned} \mathbf{L}_1 &= \sum_{i=1}^n w_{ijk} \sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} \tilde{\mathbf{U}}_{ij kts} K_{h_{\Sigma,jk}}(U_{ijt} - u) K_{h_{\Sigma,jk}}(U_{iks} - v) \frac{U_{ijt} - u}{h_{\Sigma,jk}} h_{\Sigma,jk} \frac{\partial \Sigma_{jk}}{\partial u}(u, v), \\ \mathbf{L}_2 &= \sum_{i=1}^n w_{ijk} \sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} \tilde{\mathbf{U}}_{ij kts} K_{h_{\Sigma,jk}}(U_{ijt} - u) K_{h_{\Sigma,jk}}(U_{iks} - v) \frac{U_{iks} - v}{h_{\Sigma,jk}} h_{\Sigma,jk} \frac{\partial \Sigma_{jk}}{\partial v}(u, v), \\ \mathbf{L}_3 &= \sum_{i=1}^n w_{ijk} \sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} \tilde{\mathbf{U}}_{ij kts} K_{h_{\Sigma,jk}}(U_{ijt} - u) K_{h_{\Sigma,jk}}(U_{iks} - v) \tilde{L}_{ijk}, \\ \tilde{L}_{ijk} &= \frac{1}{2} h_{\Sigma,jk}^2 \left( \frac{U_{ijt} - u}{h_{\Sigma,jk}} \right)^2 \frac{\partial^2 \Sigma_{jk}}{\partial u^2}(\delta_{ij kts1}, \delta_{ij kts2}) + \frac{1}{2} h_{\Sigma,jk}^2 \left( \frac{U_{iks} - v}{h_{\Sigma,jk}} \right)^2 \frac{\partial^2 \Sigma_{jk}}{\partial v^2}(\delta_{ij kts1}, \delta_{ij kts2}) \\ &\quad + h_{\Sigma,jk}^2 \frac{U_{ijt} - u}{h_{\Sigma,jk}} \frac{U_{iks} - v}{h_{\Sigma,jk}} \frac{\partial^2 \Sigma_{jk}}{\partial u \partial v}(\delta_{ij kts1}, \delta_{ij kts2}), \end{aligned}$$

$(\delta_{ij kts1}, \delta_{ij kts2}) \in [u - h_{\Sigma,jk}, u + h_{\Sigma,jk}] \times [v - h_{\Sigma,jk}, v + h_{\Sigma,jk}]$  and the event  $\tilde{V}_{jk} = \{(U_{ijt}, U_{iks}), t \in [T_{ij}], s \in [T_{ik}], i \in [n]\}$ . First consider  $L_1$ , which equals to the second column of  $\hat{\mathbf{\Xi}}_{jk}(u, v)$

multiplied by  $h_{\Sigma,jk} \partial \Sigma_{jk}(u, v) / \partial u$ , hence  $\mathbb{E}_U \left( \tilde{\mathbf{e}}_0^T \left[ \mathbb{E} \{ \hat{\boldsymbol{\Sigma}}_{jk}(u, v) \} \right]^{-1} \mathbf{L}_1 \right)$  equals to

$$h_{\Sigma,jk} \frac{\partial \Sigma_{jk}(u, v)}{\partial u} \tilde{\mathbf{e}}_0^T \left[ \mathbb{E} \{ \hat{\boldsymbol{\Sigma}}_{jk}(u, v) \} \right]^{-1} \mathbb{E} \{ \hat{\boldsymbol{\Sigma}}_{jk}(u, v) \} (0, 1, 0)^T = 0,$$

where  $\mathbb{E}_U$  denotes the expectation over  $\tilde{V}_{jk}$ . Following the similar procedure, we can show that

$$\mathbb{E}_U \left( \tilde{\mathbf{e}}_0^T \left[ \mathbb{E} \{ \hat{\boldsymbol{\Sigma}}_{jk}(u, v) \} \right]^{-1} \mathbf{L}_2 \right) = 0.$$

Then consider  $\mathbf{L}_3$ . By Assumption 7, we have

$$K_2 = \sup_{(j,k) \in [p]^2, (u,v) \in \mathcal{U}^2} \{ \partial^2 \Sigma_{jk}(u, v) / \partial v^2, \partial^2 \Sigma_{jk}(u, v) / \partial u^2, \partial^2 \Sigma_{jk}(u, v) / \partial u \partial v \} < \infty.$$

Each entry of  $|\mathbf{L}_3|$  is bounded by the (1, 1)th entry of  $\hat{\boldsymbol{\Sigma}}_{jk}(u, v)$  multiplied by  $2K_2 h_{\Sigma,jk}^2$ , and by  $\mathbb{E} \{ K_h(U_{ijt} - u) K_h(U_{iks} - v) \} \leq 1$ , the (1, 1)th entry of  $\mathbb{E} \{ \hat{\boldsymbol{\Sigma}}_{jk}(u, v) \}$  is bounded by 1. Note that  $\mathbb{E} \{ \hat{\boldsymbol{\Sigma}}_{jk}(u, v) \}$  is positive definite. Combining these results, we have

$$\left| \mathbb{E}_U \left( \tilde{\mathbf{e}}_0^T \left[ \mathbb{E} \{ \hat{\boldsymbol{\Sigma}}_{jk}(u, v) \} \right]^{-1} \mathbf{L}_3 \right) \right| \leq \|\mathbb{E} \{ \hat{\boldsymbol{\Sigma}}_{jk}(u, v) \}\|_{\min}^{-1} \|\mathbb{E}_U(|\mathbf{L}_3|)\| \leq cK_2 h_{\Sigma,jk}^2,$$

which implies that  $|\tilde{\Sigma}_{jk}(u, v) - \Sigma_{jk}(u, v)| \leq cK_2 h_{\Sigma,jk}^2$  for any  $(u, v) \in \mathcal{U}^2$ . Hence

$$\sup_{(u,v) \in \mathcal{U}^2} |\tilde{\Sigma}_{jk}(u, v) - \Sigma_{jk}(u, v)| = O(h_{\Sigma,jk}^2) \quad \text{and} \quad \|\tilde{\Sigma}_{jk} - \Sigma_{jk}\|_S = O(h_{\Sigma,jk}^2). \quad (\text{A.38})$$

For  $M > 0$  with the choice of  $\delta = (\log p)^{1/2} (\min_{j,k} \nu_{n,T,h,jk})^{-1/2} M \leq 1$ , it follows from the union bound of probability and (7) in Theorem 3 that

$$\begin{aligned} & \mathbb{P} \left\{ \frac{\max_{j,k \in [p]} \|\hat{\Sigma}_{jk} - \tilde{\Sigma}_{jk}\|_S}{(\log p)^{1/2} (\min_{j,k} \nu_{n,T,h,jk})^{-1/2}} \geq M \right\} \\ & \leq \sum_{j=1}^p \sum_{k=1}^p \mathbb{P} \left\{ \|\hat{\Sigma}_{jk} - \tilde{\Sigma}_{jk}\|_S \geq M (\log p)^{1/2} (\min_{j,k} \nu_{n,T,h,jk})^{-1/2} \right\} \\ & \leq \sum_{j=1}^p \sum_{k=1}^p c \exp \left( -c \nu_{n,T,h,jk} M^2 \frac{\log p}{\min_{j,k} \nu_{n,T,h,jk}} \right) \leq c \exp \{ (2 - cM^2) \log p \}. \end{aligned} \quad (\text{A.39})$$

We can choose a large  $M$  such that  $2 - cM^2 < 0$ , the right-side of (A.39) tends to 0. Hence  $\max_{j,k \in [p]} \|\hat{\Sigma}_{jk} - \tilde{\Sigma}_{jk}\|_S = O_P \{ (\log p)^{1/2} (\min_{j,k} \nu_{n,T,h,jk})^{-1/2} \}$ . Combing this with (A.38) yields that

$$\max_{j,k \in [p]} \|\hat{\Sigma}_{jk} - \Sigma_{jk}\|_S = O_P \left\{ \left( \frac{\log p}{\min_{j,k} \nu_{n,T,h,jk}} \right)^{1/2} + \max_{j,k} h_{\Sigma,jk}^2 \right\},$$

which completes the proof of (11).

The rate of convergence in (12) can be proved following a similar procedure. Let  $h_{\Sigma, \min} = \min_{j,k} h_{\Sigma,jk}$ , we can assume that  $h_{\Sigma, \min} \asymp \{ \log(p \vee n) / n \}^{\kappa_2}$  where  $\kappa_2 \in (0, 1/2]$ . Consider  $h_{\Sigma, \min}$

with some  $\kappa_2^* > 1/2$ , the corresponding rate is not faster than that with some  $\kappa_2 \in (0, 1/2]$ . Specifically, under the sparse design, the rate of  $\max_{j,k \in [p]} \sup_{(u,v) \in \mathcal{U}^2} |\widehat{\Sigma}_{jk}(u,v) - \Sigma_{jk}(u,v)|$  is  $\{\log(p \vee n)/n\}^{1/2-\kappa_2^*}$ , which is slower than  $\{\log(p \vee n)/n\}^{1/3}$  with  $\kappa_2 = 1/6$ . Under the dense design with  $\bar{T}_\Sigma \{\log(p \vee n)/n\}^{\kappa_2^*} \rightarrow 0$  and  $\bar{T}_\Sigma \{\log(p \vee n)/n\} \rightarrow 0$ , the rate is  $\{\log(p \vee n)/n\}^{1/2-\kappa_2^*} \bar{T}_\Sigma^{-1}$ , which is slower than  $\{\log(p \vee n)/n\}^{1/3} \bar{T}_\Sigma^{-2/3}$  with  $\kappa_2 \in (1/6, 1/2]$ . Under the dense design with  $\bar{T}_\Sigma \{\log(p \vee n)/n\}^{\kappa_2^*} \rightarrow 0$  and  $\bar{T}_\Sigma \{\log(p \vee n)/n\} \rightarrow \tilde{c}$  or  $\infty$ , the rate is  $\{\log(p \vee n)/n\}^{1/2-\kappa_2^*} \bar{T}_\Sigma^{-1}$ , which is slower than  $\{\log(p \vee n)/n\}^{1/2}$  with  $\kappa_2 = 1/4$ . Under the dense design with  $\bar{T}_\Sigma \{\log(p \vee n)/n\}^{\kappa_2^*} \rightarrow \tilde{c}$  or  $\infty$ , the rate is  $\{\log(p \vee n)/n\}^{1/2}$ , which is the same as  $\{\log(p \vee n)/n\}^{1/2}$  with  $\kappa_2 = 1/4$ . Based on the above four cases, if  $\kappa_2^* > 1/2$ , the corresponding rate is not faster than that with some  $\kappa_2 \in (0, 1/2]$  and hence  $\kappa_2$  that corresponds with the optimal bandwidth under sparse or dense design is in  $(0, 1/2]$ . For  $M > 0$  with the choice of  $\delta = \{\log(p \vee n)\}^{1/2} (\min_j \gamma_{n,T,h,j})^{-1/2} M \leq 1$ , by the union bound of probability and (8) in Theorem 3, we have

$$\begin{aligned}
 & \mathbb{P} \left\{ \frac{\max_{j,k \in [p]} \sup_{(u,v) \in \mathcal{U}^2} |\widehat{\Sigma}_{jk}(u,v) - \tilde{\Sigma}_{jk}(u,v)|}{\log(p \vee n)^{1/2} (\min_{j,k} \nu_{n,T,h,jk})^{-1/2}} \geq M \right\} \\
 & \leq \sum_{j=1}^p \sum_{k=1}^p \mathbb{P} \left\{ \sup_{(u,v) \in \mathcal{U}^2} |\widehat{\Sigma}_{jk}(u,v) - \tilde{\Sigma}_{jk}(u,v)| \geq M \{\log(p \vee n)\}^{1/2} (\min_{j,k} \nu_{n,T,h,jk})^{-1/2} \right\} \\
 & \leq \sum_{j=1}^p \sum_{k=1}^p \frac{cn^{\epsilon_2} \nu_{n,T,h,jk}}{h_{\Sigma, \min}^6} \exp \left\{ -c \nu_{n,T,h,jk} M^2 \frac{\log(p \vee n)}{\min_{j,k} \nu_{n,T,h,jk}} \right\} \\
 & \leq c \exp \{ (3 + \epsilon_2 + 6\kappa_2 - cM^2) \log(p \vee n) \}.
 \end{aligned}$$

We can choose a large  $M$  such that  $3 + \epsilon_2 + 6\kappa_2 - cM^2 < 0$ , the right-side of the above inequality tends to 0. Hence

$$\max_{j,k \in [p]} \sup_{(u,v) \in \mathcal{U}^2} |\widehat{\Sigma}_{jk}(u,v) - \tilde{\Sigma}_{jk}(u,v)| = O_P \left[ \left\{ \frac{\log(p \vee n)}{\min_{j,k} \nu_{n,T,h,jk}} \right\}^{1/2} \right].$$

Combing this with (A.38) yields that

$$\max_{j,k \in [p]} \sup_{(u,v) \in \mathcal{U}^2} |\widehat{\Sigma}_{jk}(u,v) - \Sigma_{jk}(u,v)| = O_P \left[ \left\{ \frac{\log(p \vee n)}{\min_{j,k} \nu_{n,T,h,jk}} \right\}^{1/2} + \max_{j,k} h_{\Sigma, jk}^2 \right],$$

which completes the proof of (12). ■

### A.5 Proof of Proposition 10

It follows from (4.43) and Lemma 4.3 of Bosq (2000) that

$$\sup_{l \in [d_j]} |\widehat{\lambda}_{jl} - \lambda_{jl}| \leq \|\widehat{\Sigma}_{jj} - \Sigma_{jj}\|_{\mathcal{S}} \quad \text{and} \quad \sup_{l \in [d_j]} \delta_{jl} \|\widehat{\phi}_{jl} - \phi_{jl}\|_2 \leq 2\sqrt{2} \|\widehat{\Sigma}_{jj} - \Sigma_{jj}\|_{\mathcal{S}}.$$

Combining the above with Theorem 7 yields the result in Proposition 10. ■

## Appendix B. Verification of the Claim in Section 5

For the mean estimator, define the set of  $r$  candidate bandwidths  $\mathcal{H}_\mu = \{h_\mu^{(1)}, \dots, h_\mu^{(r)}\}$ . In our simulations, the bandwidth for each dimension can be chosen from  $\mathcal{H}_\mu$  freely, and hence there are  $r^p$  possible outcomes. The targeted evaluation metric is  $\text{global}_{\text{opt}} = \min_{(m_1, \dots, m_p) \in [r]^p} \max_{j \in [p]} \text{MISE}(\hat{\mu}_j, h_\mu^{(m_j)})$ . We will show that  $\text{global}_{\text{opt}} = \text{MaxMISE}(\mu)$ , the right side of which is much easier to calculate as it only takes into account  $pr$  cases. On one hand, it is obvious that  $\text{global}_{\text{opt}} \leq \text{MaxMISE}(\mu)$ . On the other hand, for fixed  $(j, m_j) \in [p] \times [r]$ ,  $\text{MISE}(\hat{\mu}_j, h_\mu^{(m_j)}) \geq \min_{m \in [r]} \text{MISE}(\hat{\mu}_j, h_\mu^{(m)})$ , and hence  $\text{global}_{\text{opt}} \geq$

$$\min_{(m_1, \dots, m_p) \in [r]^p} \max_{j \in [p]} \min_{m \in [r]} \text{MISE}(\hat{\mu}_j, h_\mu^{(m)}) = \max_{j \in [p]} \min_{m \in [r]} \text{MISE}(\hat{\mu}_j, h_\mu^{(m)}) = \text{MaxMISE}(\mu).$$

Combining the above results yields  $\text{global}_{\text{opt}} = \text{MaxMISE}(\mu)$ . The corresponding claim for the covariance estimator can be verified in the same way.

## References

- Denis Bosq. *Linear Processes in Function Spaces: Theory and Applications*. Springer, New York, 2000.
- Stphane Boucheron, Gbor Lugosi, and Pascal Massart. *Concentration Inequalities: A Nonasymptotic Theory of Independence*. Oxford University Press, 2013.
- Cheng Chen, Shaojun Guo, and Xinghao Qiao. Functional linear regression: Dependence and error contamination. *Journal of Business & Economic Statistics*, 40:444–457, 2022.
- Jianqing Fan and James S. Marron. Fast implementations of nonparametric curve estimators. *Journal of Computational and Graphical Statistics*, 3:35–56, 1994.
- Yingying Fan, Natasha Foutz, Gareth M. James, and Wolfgang Jank. Functional response additive model estimation with online virtual stock markets. *The Annals of Applied Statistics*, 8:2435–2460, 2014.
- Yingying Fan, Gareth M. James, and Peter Radchenko. Functional additive regression. *The Annals of Statistics*, 43:2296–2325, 2015.
- Qin Fang, Shaojun Guo, and Xinghao Qiao. Adaptive functional thresholding for sparse covariance function estimation in high dimensions. *Journal of the American Statistical Association*, 119:1473–1485, 2024.
- Clara Happ and Sonja Greven. Multivariate functional principal component analysis for data observed on different (dimensional) domains. *Journal of the American Statistical Association*, 113:649–659, 2018.
- Xiaoyu Hu and Fang Yao. Sparse functional principal component analysis in high dimensions. *Statistica Sinica*, 32:1939–1960, 2022.
- Dehan Kong, Kaijie Xue, Fang Yao, and Hao Zhang. Partially functional linear regression in high dimensions. *Biometrika*, 103:147–159, 2016.

- Kuang-Yao Lee, Dingjue Ji, Lexin Li, Todd Constable, and Hongyu Zhao. Conditional functional graphical models. *Journal of the American Statistical Association*, 118:257–271, 2023.
- Chenlei Leng, Degui Li, Hanlin Shang, and Yingcun Xia. Covariance function estimation for high-dimensional functional time series with dual factor structures. *arXiv:2401.05784*, 2024.
- Bing Li and Eftychia Solea. A nonparametric graphical model for functional data with application to brain networks based on fMRI. *Journal of the American Statistical Association*, 113:1637–1655, 2018.
- Dong Li, Xinghao Qiao, and Zihan Wang. Factor-guided estimation of large covariance matrix function with conditional functional sparsity. *arXiv:2311.02450*, 2023.
- Yehua Li and Tailen Hsing. Uniform convergence rates for nonparametric regression and principal component analysis in functional/longitudinal data. *The Annals of Statistics*, 38:3321–3351, 2010.
- Ruiyan Luo and Xin Qi. Function-on-function linear regression by signal compression. *Journal of the American Statistical Association*, 112:690–705, 2017.
- Xinghao Qiao, Shaojun Guo, and Gareth M. James. Functional graphical models. *Journal of the American Statistical Association*, 114:211–222, 2019.
- Xinghao Qiao, Cheng Qian, Gareth M. James, and Shaojun Guo. Doubly functional graphical models in high dimensions. *Biometrika*, 107:415–431, 2020.
- Eftychia Solea and Bing Li. Copula Gaussian graphical models for functional data. *Journal of the American Statistical Association*, 117:781–793, 2022.
- Daren Wang, Zifeng Zhao, Yi Yu, and Rebecca Willett. Functional linear regression with mixed predictors. *Journal of Machine Learning Research*, 23(266):1–94, 2022.
- Kaijie Xue and Fang Yao. Hypothesis testing in large-scale functional linear regression. *Statistica Sinica*, 31:1101–1123, 2021.
- Kaijie Xue, Jin Yang, and Fang Yao. Optimal linear discriminant analysis for high-dimensional functional data. *Journal of the American Statistical Association*, 119:1055–1064, 2024.
- Fang Yao, Hans-Georg Mller, and Jane-Ling Wang. Functional data analysis for sparse longitudinal data. *Journal of the American Statistical Association*, 100:577–590, 2005a.
- Fang Yao, Hans-Georg Mller, and Jane-Ling Wang. Functional linear regression analysis for longitudinal data. *The Annals of Statistics*, 33:2873–2903, 2005b.
- Javier Zapata, Sang-Yun Oh, and Alexander Petersen. Partial separability and functional graphical models for multivariate gaussian processes. *Biometrika*, 109:665–681, 2022.

Jin-Ting Zhang and Jianwei Chen. Statistical inferences for functional data. *The Annals of Statistics*, 35:1052–1079, 2007.

Xiaoke Zhang and Jane-Ling Wang. From sparse to dense functional data and beyond. *The Annals of Statistics*, 44:2281–2321, 2016.

Boxin Zhao, Y. Samuel Wang, and Mladen Kolar. FuDGE: A method to estimate a functional differential graph in a high-dimensional setting. *Journal of Machine Learning Research*, 23(82):1–82, 2022.

Hongxiao Zhu, Nate Strawn, and David B. Dunson. Bayesian graphical models for multivariate functional data. *Journal of Machine Learning Research*, 17(204):1–27, 2016.